

Measurement of the W boson mass

B. Abbott,³⁰ M. Abolins,²⁷ B. S. Acharya,⁴⁵ I. Adam,¹² D. L. Adams,³⁹ M. Adams,¹⁷ S. Ahn,¹⁴ H. Aihara,²³ G. A. Alves,¹⁰ N. Amos,²⁶ E. W. Anderson,¹⁹ R. Astur,⁴⁴ M. M. Baarmand,⁴⁴ A. Baden,²⁵ V. Balamurali,³⁴ J. Balderston,¹⁶ B. Baldin,¹⁴ S. Banerjee,⁴⁵ J. Bantly,⁵ E. Barberis,²³ J. F. Bartlett,¹⁴ K. Bazizi,⁴¹ A. Belyaev,²⁸ S. B. Beri,³⁶ I. Bertram,³³ V. A. Bezzubov,³⁷ P. C. Bhat,¹⁴ V. Bhatnagar,³⁶ M. Bhattacharjee,⁴⁴ N. Biswas,³⁴ G. Blazey,³² S. Blessing,¹⁵ P. Bloom,⁷ A. Boehnlein,¹⁴ N. I. Bojko,³⁷ F. Borchering,¹⁴ C. Boswell,⁹ A. Brandt,¹⁴ R. Brock,²⁷ A. Bross,¹⁴ D. Buchholz,³³ V. S. Burtovoi,³⁷ J. M. Butler,³ W. Carvalho,¹⁰ D. Casey,⁴¹ Z. Casilum,⁴⁴ H. Castilla-Valdez,¹¹ D. Chakraborty,⁴⁴ S.-M. Chang,³¹ S. V. Chekulaev,³⁷ L.-P. Chen,²³ W. Chen,⁴⁴ S. Choi,⁴³ S. Chopra,²⁶ B. C. Choudhary,⁹ J. H. Christenson,¹⁴ M. Chung,¹⁷ D. Claes,²⁹ A. R. Clark,²³ W. G. Cobau,²⁵ J. Cochran,⁹ L. Coney,³⁴ W. E. Cooper,¹⁴ C. Cretsinger,⁴¹ D. Cullen-Vidal,⁵ M. A. C. Cummings,³² D. Cutts,⁵ O. I. Dahl,²³ K. Davis,² K. De,⁴⁶ K. Del Signore,²⁶ M. Demarteau,¹⁴ D. Denisov,¹⁴ S. P. Denisov,³⁷ H. T. Diehl,¹⁴ M. Diesburg,¹⁴ G. Di Loreto,²⁷ P. Draper,⁴⁶ Y. Ducros,⁴² L. V. Dudko,²⁸ S. R. Dugad,⁴⁵ D. Edmunds,²⁷ J. Ellison,⁹ V. D. Elvira,⁴⁴ R. Engelmann,⁴⁴ S. Eno,²⁵ G. Eppley,³⁹ P. Ermolov,²⁸ O. V. Eroshin,³⁷ V. N. Evdokimov,³⁷ T. Fahland,⁸ M. K. Fatyga,⁴¹ S. Feher,¹⁴ D. Fein,² T. Ferbel,⁴¹ G. Finocchiaro,⁴⁴ H. E. Fisk,¹⁴ Y. Fisyak,⁷ E. Flattum,¹⁴ G. E. Forden,² M. Fortner,³² K. C. Frame,²⁷ S. Fuess,¹⁴ E. Gallas,⁴⁶ A. N. Galyaev,³⁷ P. Gartung,⁹ T. L. Geld,²⁷ R. J. Genik II,²⁷ K. Genser,¹⁴ C. E. Gerber,¹⁴ B. Gibbard,⁴ S. Glenn,⁷ B. Gobbi,³³ A. Goldschmidt,²³ B. Gómez,¹ G. Gómez,²⁵ P. I. Goncharov,³⁷ J. L. González Solís,¹¹ H. Gordon,⁴ L. T. Goss,⁴⁷ K. Gounder,⁹ A. Goussiou,⁴⁴ N. Graf,⁴ P. D. Grannis,⁴⁴ D. R. Green,¹⁴ H. Greenlee,¹⁴ G. Grim,⁷ S. Grinstein,⁶ N. Grossman,¹⁴ P. Grudberg,²³ S. Grünendahl,¹⁴ G. Guglielmo,³⁵ J. A. Guida,² J. M. Guida,⁵ A. Gupta,⁴⁵ S. N. Gurzhiev,³⁷ P. Gutierrez,³⁵ Y. E. Gutnikov,³⁷ N. J. Hadley,²⁵ H. Haggerty,¹⁴ S. Hagopian,¹⁵ V. Hagopian,¹⁵ K. S. Hahn,⁴¹ R. E. Hall,⁸ P. Hanlet,³¹ S. Hansen,¹⁴ J. M. Hauptman,¹⁹ D. Hedin,³² A. P. Heinson,⁹ U. Heintz,¹⁴ R. Hernández-Montoya,¹¹ T. Heuring,¹⁵ R. Hirosky,¹⁷ J. D. Hobbs,¹⁴ B. Hoeneisen,¹ J. S. Hoftun,⁵ F. Hsieh,²⁶ Ting Hu,⁴⁴ Tong Hu,¹⁸ T. Huehn,⁹ A. S. Ito,¹⁴ E. James,² J. Jaques,³⁴ S. A. Jerger,²⁷ R. Jesik,¹⁸ J. Z.-Y. Jiang,⁴⁴ T. Joffe-Minor,³³ K. Johns,² M. Johnson,¹⁴ A. Jonckheere,¹⁴ M. Jones,¹⁶ H. Jöstlein,¹⁴ S. Y. Jun,³³ C. K. Jung,⁴⁴ S. Kahn,⁴ G. Kalbfleisch,³⁵ J. S. Kang,²⁰ D. Karmanov,²⁸ D. Karmgard,¹⁵ R. Kehoe,³⁴ M. L. Kelly,³⁴ C. L. Kim,²⁰ S. K. Kim,⁴³ A. Klatchko,¹⁵ B. Klima,¹⁴ C. Klopfenstein,⁷ V. I. Klyukhin,³⁷ V. I. Kochetkov,³⁷ J. M. Kohli,³⁶ D. Koltick,³⁸ A. V. Kostritskiy,³⁷ J. Kotcher,⁴ A. V. Kotwal,¹² J. Kourlas,³⁰ A. V. Kozelov,³⁷ E. A. Kozlovski,³⁷ J. Krane,²⁹ M. R. Krishnaswamy,⁴⁵ S. Krzywdzinski,¹⁴ S. Kunori,²⁵ S. Lami,⁴⁴ R. Lander,⁷ F. Landry,²⁷ G. Landsberg,¹⁴ B. Lauer,¹⁹ A. Leflat,²⁸ H. Li,⁴⁴ J. Li,⁴⁶ Q. Z. Li-Demarteau,¹⁴ J. G. R. Lima,⁴⁰ D. Lincoln,²⁶ S. L. Linn,¹⁵ J. Linnemann,²⁷ R. Lipton,¹⁴ Y. C. Liu,³³ F. Lobkowicz,⁴¹ S. C. Loken,²³ S. Lökös,⁴⁴ L. Lueking,¹⁴ A. L. Lyon,²⁵ A. K. A. Maciel,¹⁰ R. J. Madaras,²³ R. Madden,¹⁵ L. Magaña-Mendoza,¹¹ V. Manankov,²⁸ S. Mani,⁷ H. S. Mao,¹⁴ R. Markeloff,³² T. Marshall,¹⁸ M. I. Martin,¹⁴ K. M. Mauritz,¹⁹ B. May,³³ A. A. Mayorov,³⁷ R. McCarthy,⁴⁴ J. McDonald,¹⁵ T. McKibben,¹⁷ J. McKinley,²⁷ T. McMahon,³⁵ H. L. Melanson,¹⁴ M. Merkin,²⁸ K. W. Merritt,¹⁴ H. Miettinen,³⁹ A. Mincer,³⁰ C. S. Mishra,¹⁴ N. Mokhov,¹⁴ N. K. Mondal,⁴⁵ H. E. Montgomery,¹⁴ P. Mooney,¹ H. da Motta,¹⁰ C. Murphy,¹⁷ F. Nang,² M. Narain,¹⁴ V. S. Narasimham,⁴⁵ A. Narayanan,² H. A. Neal,²⁶ J. P. Negret,¹ P. Nemethy,³⁰ D. Norman,⁴⁷ L. Oesch,²⁶ V. Oguri,⁴⁰ E. Oliveira,¹⁰ E. Oltman,²³ N. Oshima,¹⁴ D. Owen,²⁷ P. Padley,³⁹ A. Para,¹⁴ Y. M. Park,²¹ R. Partridge,⁵ N. Parua,⁴⁵ M. Paterno,⁴¹ B. Pawlik,²² J. Perkins,⁴⁶ M. Peters,¹⁶ R. Piegaia,⁶ H. Piekarz,¹⁵ Y. Pischalnikov,³⁸ V. M. Podstavkov,³⁷ B. G. Pope,²⁷ H. B. Prosper,¹⁵ S. Protopopescu,⁴ J. Qian,²⁶ P. Z. Quintas,¹⁴ R. Raja,¹⁴ S. Rajagopalan,⁴ O. Ramirez,¹⁷ L. Rasmussen,⁴⁴ S. Reucroft,³¹ M. Rijssenbeek,⁴⁴ T. Rockwell,²⁷ M. Roco,¹⁴ N. A. Roe,²³ P. Rubinov,³³ R. Ruchti,³⁴ J. Rutherford,² A. Sánchez-Hernández,¹¹ A. Santoro,¹⁰ L. Sawyer,²⁴ R. D. Schamberger,⁴⁴ H. Schellman,³³ J. Sculli,³⁰ E. Shabalina,²⁸ C. Shaffer,¹⁵ H. C. Shankar,⁴⁵ R. K. Shivpuri,¹³ M. Shupe,² H. Singh,⁹ J. B. Singh,³⁶ V. Sirotenko,³² W. Smart,¹⁴ E. Smith,³⁵ R. P. Smith,¹⁴ R. Snihur,³³ G. R. Snow,²⁹ J. Snow,³⁵ S. Snyder,⁴ J. Solomon,¹⁷ P. M. Sood,³⁶ M. Sosebee,⁴⁶ N. Sotnikova,²⁸ M. Souza,¹⁰ A. L. Spadafora,²³ G. Steinbrück,³⁵ R. W. Stephens,⁴⁶ M. L. Stevenson,²³ D. Stewart,²⁶ F. Stichelbaut,⁴⁴ D. A. Stoianova,³⁷ D. Stoker,⁸ M. Strauss,³⁵ K. Streets,³⁰ M. Strovink,²³ A. Sznajder,¹⁰ P. Tamburello,²⁵ J. Tarazi,⁸ M. Tartaglia,¹⁴ T. L. T. Thomas,³³ J. Thompson,²⁵ T. G. Trippe,²³ P. M. Tuts,¹² N. Varelas,¹⁷ E. W. Varnes,²³ D. Vittoe,² A. A. Volkov,³⁷ A. P. Vorobiev,³⁷ H. D. Wahl,¹⁵ G. Wang,¹⁵ J. Warchol,³⁴ G. Watts,⁵ M. Wayne,³⁴ H. Weerts,²⁷ A. White,⁴⁶ J. T. White,⁴⁷ J. A. Wightman,¹⁹ S. Willis,³² S. J. Wimpenny,⁹ J. V. D. Wirjawan,⁴⁷ J. Womersley,¹⁴ E. Won,⁴¹ D. R. Wood,³¹ H. Xu,⁵ R. Yamada,¹⁴ P. Yamin,⁴ J. Yang,³⁰ T. Yasuda,³¹ P. Yepes,³⁹ C. Yoshikawa,¹⁶ S. Youssef,¹⁵ J. Yu,¹⁴ Y. Yu,⁴³ Z. H. Zhu,⁴¹ D. Zieminska,¹⁸ A. Zieminski,¹⁸ E. G. Zverev,²⁸ and A. Zylberstejn⁴²

¹Universidad de los Andes, Bogotá, Colombia²University of Arizona, Tucson, Arizona 85721³Boston University, Boston, Massachusetts 02215⁴Brookhaven National Laboratory, Upton, New York 11973⁵Brown University, Providence, Rhode Island 02912⁶Universidad de Buenos Aires, Buenos Aires, Argentina⁷University of California, Davis, California 95616⁸University of California, Irvine, California 92697⁹University of California, Riverside, California 92521

- ¹⁰LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil
¹¹CINVESTAV, Mexico City, Mexico
¹²Columbia University, New York, New York 10027
¹³Delhi University, Delhi, India 110007
¹⁴Fermi National Accelerator Laboratory, Batavia, Illinois 60510
¹⁵Florida State University, Tallahassee, Florida 32306
¹⁶University of Hawaii, Honolulu, Hawaii 96822
¹⁷University of Illinois at Chicago, Chicago, Illinois 60607
¹⁸Indiana University, Bloomington, Indiana 47405
¹⁹Iowa State University, Ames, Iowa 50011
²⁰Korea University, Seoul, Korea
²¹Kyungshung University, Pusan, Korea
²²Institute of Nuclear Physics, Kraków, Poland
²³Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720
²⁴Louisiana Tech University, Ruston, Louisiana 71272
²⁵University of Maryland, College Park, Maryland 20742
²⁶University of Michigan, Ann Arbor, Michigan 48109
²⁷Michigan State University, East Lansing, Michigan 48824
²⁸Moscow State University, Moscow, Russia
²⁹University of Nebraska, Lincoln, Nebraska 68588
³⁰New York University, New York, New York 10003
³¹Northeastern University, Boston, Massachusetts 02115
³²Northern Illinois University, DeKalb, Illinois 60115
³³Northwestern University, Evanston, Illinois 60208
³⁴University of Notre Dame, Notre Dame, Indiana 46556
³⁵University of Oklahoma, Norman, Oklahoma 73019
³⁶University of Panjab, Chandigarh 16-00-14, India
³⁷Institute for High Energy Physics, 142-284 Protvino, Russia
³⁸Purdue University, West Lafayette, Indiana 47907
³⁹Rice University, Houston, Texas 77005
⁴⁰Universidade do Estado do Rio de Janeiro, Brazil
⁴¹University of Rochester, Rochester, New York 14627
⁴²CEA, DAPNIA/Service de Physique des Particules, CE-SACLAY, Gif-sur-Yvette, France
⁴³Seoul National University, Seoul, Korea
⁴⁴State University of New York, Stony Brook, New York 11794
⁴⁵Tata Institute of Fundamental Research, Colaba, Mumbai 400005, India
⁴⁶University of Texas, Arlington, Texas 76019
⁴⁷Texas A&M University, College Station, Texas 77843

(D0 Collaboration)

(Received 23 December 1997; published 24 September 1998)

We present a measurement of the W boson mass using data collected by the D0 experiment at the Fermilab Tevatron during 1994–1995. We identify W bosons by their decays to $e\nu$ final states. We extract the W mass M_W by fitting the transverse mass and transverse electron momentum spectra from a sample of 28 323 $W \rightarrow e\nu$ decay candidates. We use a sample of 3563 dielectron events, mostly due to $Z \rightarrow ee$ decays, to constrain our model of the detector response. From the transverse mass fit we measure $M_W = 80.44 \pm 0.10(\text{stat}) \pm 0.07(\text{syst})$ GeV. Combining this with our previously published result from data taken in 1992–1993, we obtain $M_W = 80.43 \pm 0.11$ GeV. [S0556-2821(98)03519-X]

PACS number(s): 14.70.Fm, 12.15.Ji, 13.38.Be, 13.85.Qk

I. INTRODUCTION

In this article we describe the most precise measurement to date of the mass of the W boson, using data collected in 1994–1995 with the D0 detector at the Fermilab Tevatron $p\bar{p}$ collider [1–3].

The study of the properties of the W boson began in 1983 with its discovery by the UA1 [4] and UA2 [5] Collabora-

tions at the CERN $p\bar{p}$ collider. Together with the discovery of the Z boson in the same year [6,7], it provided direct confirmation of the unified model of weak and electromagnetic interactions [8], which—together with QCD—is now called the standard model.

Since the W and Z bosons are carriers of the weak force, their properties are intimately coupled to the structure of the model. The properties of the Z boson have been studied in

FIG. 1. Loop diagrams contributing to the W boson mass.

great detail in e^+e^- collisions [9]. The study of the W boson has proved to be significantly more difficult, since it is charged and therefore cannot be resonantly produced in e^+e^- collisions. Until recently its direct study has therefore been the realm of experiments at $p\bar{p}$ colliders, which have performed the most precise direct measurements of the W boson mass [10–12]. Direct measurements of the W boson mass have also been carried out at the CERN e^+e^- collider LEP2 [13–16] using nonresonant W pair production. A summary of these measurements can be found in Table XV at the end of this article.

The standard model links the W boson mass to other parameters:

$$M_W = \left(\frac{\pi\alpha}{\sqrt{2}G_F} \right)^{1/2} \frac{1}{\sin\theta_W\sqrt{1-\Delta r}}. \quad (1)$$

in the ‘‘on-shell’’ scheme [17],

$$\cos\theta_W = \frac{M_W}{M_Z}, \quad (2)$$

where θ_W is the weak mixing angle. Aside from the radiative corrections Δr , the W boson mass is thus determined by three precisely measured quantities, the mass of the Z boson M_Z [9], the Fermi constant G_F [18], and the electromagnetic coupling constant α evaluated at $Q^2 = M_Z^2$ [19]:


$$M_Z = 91.1867 \pm 0.0020 \text{ GeV}, \quad (3)$$

$$G_F = (1.16639 \pm 0.00002) \times 10^{-5} \text{ GeV}^{-2}, \quad (4)$$

$$\alpha = (128.896 \pm 0.090)^{-1}. \quad (5)$$

From the measured W boson mass, we can derive the size of the radiative corrections Δr . Within the framework of the standard model, these corrections are dominated by loops involving the top quark and the Higgs boson (see Fig. 1). The correction from the $t\bar{b}$ loop is substantial because of the large mass difference between the two quarks. It is proportional to m_t^2 for large values of the top quark mass m_t . Since m_t has been measured [20], this contribution can be calculated within the standard model. For a large Higgs boson mass m_H , the correction from the Higgs loop is proportional to $\ln m_H$. In extensions to the standard model, new particles may give rise to additional corrections to the value of M_W . In the minimal supersymmetric extension of the standard model (MSSM), for example, additional corrections can increase the predicted W mass by up to 250 MeV [21].

A measurement of the W boson mass therefore constitutes a test of the standard model. In conjunction with a measurement of the top quark mass, the standard model predicts M_W

FIG. 2. Lowest-order diagrams for W and Z boson production.

up to a 200 MeV uncertainty due to the unknown Higgs boson mass. By comparing with the measured value of the W boson mass, we can constrain the mass of the Higgs boson, the agent of the electroweak symmetry breaking that has up to now eluded experimental detection. A discrepancy with the range allowed by the standard model could indicate new physics. The experimental challenge is thus to measure the W boson mass to sufficient precision, about 0.1%, to be sensitive to these corrections.

II. OVERVIEW

A. Conventions

We use a Cartesian coordinate system with the z axis defined by the direction of the proton beam, the x axis pointing radially out of the Tevatron ring, and the y axis pointing up. A vector \vec{p} is then defined in terms of its projections on these three axes, p_x , p_y , and p_z . Since protons and antiprotons in the Tevatron are unpolarized, all physical processes are invariant with respect to rotations around the beam direction. It is therefore convenient to use a cylindrical coordinate system, in which the same vector is given by the magnitude of its component transverse to the beam direction, p_T , its azimuth ϕ , and p_z . In $p\bar{p}$ collisions the center-of-mass frame of the parton-parton collisions is approximately at rest in the plane transverse to the beam direction, but has an undetermined motion along the beam direction. Therefore the plane transverse to the beam direction is of special importance, and sometimes we work with two-dimensional vectors defined in the x - y plane. They are written with a subscript T , e.g., \vec{p}_T . We also use spherical coordinates by replacing p_z with the colatitude θ or the pseudorapidity $\eta = -\ln \tan(\theta/2)$. The origin of the coordinate system is in general the reconstructed position of the $p\bar{p}$ interaction when describing the interaction and the geometrical center of the detector when describing the detector. For convenience, we use units in which $c = \hbar = 1$.

B. W and Z boson production and decay

In $p\bar{p}$ collisions at $\sqrt{s} = 1.8 \text{ TeV}$, W and Z bosons are produced predominantly through quark-antiquark annihilation. Figure 2 shows the lowest-order diagrams. The quarks in the initial state may radiate gluons, which are usually very soft, but may sometimes be energetic enough to give rise to hadron jets in the detector. In the reaction the initial proton and antiproton break up and the fragments hadronize. We refer to everything except the vector boson and its decay products collectively as the underlying event. Since the initial proton and antiproton momentum vectors add to zero, the

same must be true for the vector sum of all final-state momenta, and therefore the vector boson recoils against all particles in the underlying event. The sum of the transverse momenta of the recoiling particles must balance the transverse momentum of the boson, which is typically small compared to its mass, but has a long tail to large values.

We identify W and Z bosons by their leptonic decays. The D0 detector (Sec. III) is best suited for a precision measurement of electrons and positrons,¹ and we therefore use the decay channel $W \rightarrow e\nu$ to measure the W boson mass. $Z \rightarrow ee$ decays serve as an important calibration sample. About 11% of the W bosons decay to $e\nu$, and about 3.3% of the Z bosons decay to ee . The leptons typically have transverse momenta of about half the mass of the decaying boson and are well isolated from other large energy deposits in the calorimeter. Intermediate vector boson decays are the dominant source of isolated high- p_T leptons at the Tevatron, and therefore these decays allow us to select a clean sample of W and Z boson decays.

C. Event characteristics

In events due to the process $p\bar{p} \rightarrow (W \rightarrow e\nu) + X$, where X stands for the underlying event, we detect the electron and all particles recoiling against the W with pseudorapidity $-4 < \eta < 4$. The neutrino escapes undetected. In the calorimeter we cannot resolve individual recoil particles, but we measure their energies summed over detector segments. Recoil particles with $|\eta| > 4$ escape unmeasured through the beam pipe, possibly carrying away substantial momentum along the beam direction. This means that we cannot measure the sum of the z components of the recoil momenta, u_z , precisely. Since these particles escape at a very small angle with respect to the beam, their transverse momenta are typically small and can be neglected in the sum of the transverse recoil momenta, \vec{u}_T . We measure \vec{u}_T by summing the observed energy flow vectorially over all detector segments. Thus we reduce the reconstruction of every candidate event to a measurement of the electron momentum $\vec{p}(e)$ and \vec{u}_T .

Since the neutrino escapes undetected, the sum of all measured final-state transverse momenta does not add to zero. The missing transverse momentum \vec{p}_T , required to balance the transverse momentum sum, is a measure of the transverse momentum of the neutrino. The neutrino momentum component along the beam direction cannot be determined, because u_z is not measured well. The signature of a $W \rightarrow e\nu$ decay is therefore an isolated high- p_T electron and large missing transverse momentum.

In the case of $Z \rightarrow ee$ decays the signature consists of two isolated high- p_T electrons and we measure the momenta of both leptons, $\vec{p}(e_1)$ and $\vec{p}(e_2)$, and \vec{u}_T in the detector.

D. Mass measurement strategy

Since $p_z(\nu)$ is unknown, we cannot reconstruct the $e\nu$ invariant mass for $W \rightarrow e\nu$ candidate events and therefore must resort to other kinematic variables for the mass measurement.

¹In the following we use “electron” generically for both electrons and positrons.




FIG. 3. m_T spectrum for W bosons with $q_T=0$ (solid line), with the correct q_T distribution (\bullet), and with detector resolutions (shaded area).

For recent measurements [10–12] the transverse mass

$$m_T = \sqrt{2p_T(e)p_T(\nu)\{1 - \cos[\phi(e) - \phi(\nu)]\}} \quad (6)$$

was used. This variable has the advantage that its spectrum is relatively insensitive to the production dynamics of the W . Corrections to m_T due to the motion of the W are of order $(q_T/M_W)^2$, where q_T is the transverse momentum of the W boson. It is also insensitive to selection biases that prefer certain event topologies (Sec. VI C). However, it makes use of the inferred neutrino p_T and is therefore sensitive to the response of the detector to the recoil particles.

The electron p_T spectrum provides an alternative measurement of the W mass. It is measured with better resolution than the neutrino p_T and is insensitive to the recoil momentum measurement. However, its shape is sensitive to the motion of the W and receives corrections of order q_T/M_W . It thus requires a better understanding of the W boson production dynamics than the m_T spectrum.

The m_T and $p_T(e)$ spectra thus provide us with two complementary measurements. This is illustrated in Figs. 3 and 4, which show the effect of the motion of the W bosons and the detector resolutions on the shape of each of the two spectra. The solid line shows the shape of the distribution before the detector simulation and with $q_T=0$. The points show the shape after q_T is added to the system, and the shaded histogram also includes the detector simulation. We observe that the shape of the m_T spectrum is dominated by detector resolutions and the shape of the $p_T(e)$ spectrum by the motion of the W . By performing the measurement using both spectra, we provide a powerful cross-check with complementary systematics.

Both spectra are equally sensitive to the electron energy response of the detector. We calibrate this response by forcing the observed dielectron mass peak in the $Z \rightarrow ee$ sample to agree with the known Z mass [9] (Sec. VI). This means that we effectively measure the ratio of W and Z masses, which is equivalent to a measurement of the W mass because the Z mass is known precisely.




FIG. 4. $p_T(e)$ spectrum for W bosons with $q_T=0$ (solid line), with the correct q_T distribution (\bullet), and with detector resolutions (shaded area).

To carry out these measurements we perform a maximum likelihood fit to the spectra. Since the shape of the spectra, including all the experimental effects, cannot be computed analytically, we need a Monte Carlo simulation program that can predict the shape of the spectra as a function of the W mass. To perform a measurement of the W mass to a precision of order 100 MeV, we have to estimate individual systematic effects to 10 MeV. This requires a Monte Carlo sample of 2.5×10^6 accepted W bosons for each such effect. The program therefore must be capable of generating large samples in a reasonable time. We achieve the required performance by employing a parametrized model of the detector response.

We next summarize the aspects of the accelerator and detector that are important for our measurement (Sec. III). Then we describe the data selection (Sec. IV) and the fast Monte Carlo model (Sec. V). Most parameters in the model are determined from our data. We describe the determination of the various components of the Monte Carlo model in Secs. VI–IX. After tuning the model we fit the kinematic spectra (Sec. X), perform some consistency checks (Sec. XI), and discuss the systematic uncertainties (Sec. XII). Section XIII summarizes the results and presents the conclusions.

III. EXPERIMENTAL SETUP

A. Accelerator

The Fermilab Tevatron [22] collides proton and antiproton beams at a center-of-mass energy of $\sqrt{s}=1.8$ TeV. Six bunches each of protons and antiprotons circulate around the ring in opposite directions. Bunches cross at the intersection regions every $3.5 \mu\text{s}$. During the 1994–1995 running period, the accelerator reached a peak luminosity of $2.5 \times 10^{31} \text{ cm}^{-2} \text{ s}^{-1}$ and delivered an integrated luminosity of about 100 pb^{-1} .

The Tevatron tunnel also houses a 150 GeV proton synchrotron, called the Main Ring, which is used as an injector for the Tevatron. The Main Ring also serves to accelerate




FIG. 5. Cutaway view of the D0 calorimeter and tracking system.

protons for antiproton production during collider operation. Since the Main Ring beam pipe passes through the outer section of the D0 calorimeter, passing proton bunches give rise to backgrounds in the detector. We eliminate this background using timing cuts based on the accelerator clock signal.

B. Detector

1. Overview

The D0 detector consists of three major subsystems: a central detector, a calorimeter (Fig. 5), and a muon spectrometer. It is described in detail in Ref. [23]. We describe only the features that are most important for this measurement.

2. Central detector

The central detector is designed to measure the trajectories of charged particles. It consists of a vertex drift chamber, a transition radiation detector, a central drift chamber (CDC), and two forward drift chambers (FDCs). There is no central magnetic field. The CDC covers the region $|\eta| < 1.0$. It is a jet-type drift chamber with delay lines to give the hit coordinates in the r - z plane. The (FDCs) cover the region $1.4 < |\eta| < 3.0$.

3. Calorimeter

The calorimeter is the most important part of the detector for this measurement. It is a sampling calorimeter and uses uranium absorber plates and liquid argon as the active medium. It is divided into three parts: a central calorimeter (CC) and two end calorimeters (ECs), each housed in its own cryostat. Each is segmented into an electromagnetic (EM) section, a fine hadronic (FH) section, and a coarse hadronic (CH) section, with increasingly coarser sampling. The CC-EM section is constructed of 32 azimuthal modules. The entire calorimeter is divided into about 5000 pseudoprojective towers, each covering 0.1×0.1 in $\eta \times \phi$. The EM section is segmented into four layers, 2, 2, 7, and 10 radiation lengths thick. The third layer, in which electromagnetic




FIG. 6. Pedestal spectrum of a central calorimeter cell, where the mean pedestal has been subtracted. The shaded region is the events removed by the zero suppression.

showers typically reach their maximum, is transversely segmented into cells covering 0.05×0.05 in $\eta \times \phi$. The hadronic section is segmented into four layers (CC) or five layers (EC). The entire calorimeter is 7–9 nuclear interaction lengths thick. There are no projective cracks in the calorimeter, and it provides hermetic and almost uniform coverage for particles with $|\eta| < 4$. Figure 5 shows a view of the calorimeter and the central detector.

The signals from arrays of 2×2 calorimeter towers, covering 0.2×0.2 in $\eta \times \phi$, are added together electronically for the EM section only and for all sections, and shaped with a fast rise time for use in the level 1 trigger. We refer to these arrays of 2×2 calorimeter towers as “trigger towers.”

Figure 6 shows the pedestal spectrum of a calorimeter cell. The spectrum has an asymmetric tail from ionization caused by the intrinsic radioactivity of the uranium absorber plates. The data are corrected such that the mean pedestal is zero for each cell. To reduce the amount of data that have to be stored, the calorimeter readout is zero suppressed. Only cells with a signal that deviates from zero by more than twice




FIG. 7. Response of the liquid argon in the central calorimeter as monitored by α and β sources.




FIG. 8. Percentage change in the central calorimeter gains over the course of the run.

the rms of the pedestal distribution are read out. This region of the pedestal spectrum is indicated by the shaded region in Fig. 6. Because of its asymmetry, the spectrum does not average to zero after zero suppression. Thus the zero suppression effectively causes a pedestal shift.

The liquid argon has unit gain, and therefore the calorimeter response was extremely stable during the entire run. Figure 7 shows the response of the liquid argon as monitored with radioactive sources of α and β particles. Figures 8 and 9 show the gains and pedestals of a typical readout channel throughout the run.

The EM calorimeter provides a measurement of energy and position of the electrons from the W and Z decays. Because of the fine segmentation of the third layer, we can measure the position of the shower centroid with a precision of 2.5 mm in the azimuthal direction and 1 cm in the z direction.

We study the response of the EM calorimeter to electrons in beam tests [24]. To reconstruct the electron energy we add the signals a_i observed in each EM layer ($i = 1, \dots, 4$) and the first FH layer ($i = 5$) of an array of 5×5 calorimeter towers, centered on the most energetic tower, weighted by a layer dependent sampling weight s_i :

$$E = A \sum_{i=1}^5 s_i a_i - \delta_{EM}. \quad (7)$$

To determine the sampling weights, we minimize




FIG. 9. Change in the central calorimeter pedestals over the course of the run.




FIG. 10. Fractional deviation of the reconstructed electron energy from the beam momentum from beam tests of a CC-EM module.

$$\chi^2 = \sum \frac{(p_{\text{beam}} - E)^2}{\sigma_{\text{EM}}^2}, \quad (8)$$

where the sum runs over all events and σ_{EM} is the resolution given in Eq. (9). We obtain $A = 2.96$ MeV/analogue to digital converter (ADC) count, $\delta_{\text{EM}} = -347$ MeV, $s_1 = 1.31$, $s_2 = 0.85$, $s_4 = 0.98$, and $s_5 = 1.84$. We arbitrarily fix $s_3 = 1$. The value of δ_{EM} depends on the amount of dead material in front of the calorimeter. The parameters s_1 – s_4 weight the four EM layers and s_5 the first FH layer. Figure 10 shows the fractional deviation of E as a function of the beam momentum p_{beam} . Above 10 GeV they deviate by less than 0.3% from each other.

The fractional energy resolution can be parametrized as a function of electron energy using constant, sampling, and noise terms as

$$\left(\frac{\sigma_{\text{EM}}}{E}\right)^2 = c_{\text{EM}}^2 + \left(\frac{s_{\text{EM}}}{\sqrt{E} \sin \theta}\right)^2 + \left(\frac{n_{\text{EM}}}{E}\right)^2, \quad (9)$$

with $c_{\text{EM}} = 0.003$, $s_{\text{EM}} = 0.135$ GeV^{1/2} [25,26], and $n_{\text{EM}} = 0.43$ GeV in the central calorimeter. The angle θ is the colatitude of the electron. Figure 11 shows the fractional electron energy resolution versus beam momentum for a CC-EM module. The line shows the parametrization of the resolution from Eq. (9).

4. Luminosity monitor

Two arrays of scintillator hodoscopes, mounted in front of the EC cryostats, register hits with a 220 ps time resolution. They serve to detect that an inelastic $p\bar{p}$ interaction has taken place. The particles from the breakup of the proton give rise to hits in the hodoscopes on one side of the detector that are tightly clustered in time. The detector has a 91% acceptance for inelastic $p\bar{p}$ interactions. For events with a single interaction, the location of the interaction vertex can be deter-




FIG. 11. Fractional electron energy resolution measured in beam tests of a CC-EM module for the data (●) and the parametrization (solid line).

mined with a resolution of 3 cm from the time difference between the hits on the two sides of the detector for use in the level 2 trigger. This array is also called the level 0 trigger because the detection of an inelastic $p\bar{p}$ interaction is a basic requirement of most trigger conditions.

5. Trigger

Readout of the detector is controlled by a two-level trigger system.

Level 1 consists of an and/or network, which can be programmed to trigger on a $p\bar{p}$ crossing in a number of preselected conditions are true. The level 1 trigger decision is taken within the 3.5 μs time interval between crossings. As an extension to level 1, a trigger processor (level 1.5) may be invoked to execute simple algorithms on the limited information available at the time of a level 1 accept. For electrons, the processor uses the energy deposits in each trigger tower as inputs. The detector cannot accept any triggers until the level 1.5 processor completes execution and accepts or rejects the event.

Level 2 of the trigger consists of a farm of 48 VAXstation 4000's. At this level the complete event is available. More sophisticated algorithms refine the trigger decisions, and events are accepted based on preprogrammed conditions. Events accepted by level 2 are written to magnetic tape for off-line reconstruction.

IV. DATA SELECTION

A. Trigger

The conditions required at trigger level 1 for W and Z candidates are the following.

$p\bar{p}$ interaction. Level 0 hodoscopes register hits consistent with a $p\bar{p}$ interaction. This condition accepts 98.6% of all W and Z bosons produced.

Main Ring veto. No Main Ring proton bunch passes through the detector less than 800 ns before or after the $p\bar{p}$ crossing, and no protons were injected into the Main Ring




FIG. 12. Relative efficiency of the level 2 electron filter for a threshold of 20 GeV. The arrow indicates the cut applied in the final event selection.

less than 400 ms before the $p\bar{p}$ crossing.

EM trigger towers. There are one or more EM trigger towers with $E \sin \theta > T$, where E is the energy measured in the tower, θ its angle with the beam measured from the center of the detector, and T a programmable threshold. This requirement is fully efficient for electrons with $p_T > 2T$.

The level 1.5 processor recomputes the transverse electron energy by adding the adjacent EM trigger tower with the largest signal to the EM trigger tower that exceeded the level 1 threshold. In addition, the signal in the EM trigger tower that exceeded the level 1 threshold must constitute at least 85% of the signal registered in this tower if the hadronic layers are also included. This EM fraction requirement is fully efficient for electron candidates that pass our off-line selection (Sec. IV D).

Level 2 uses the EM trigger tower that exceeded the level 1 threshold as a starting point. The level 2 algorithm finds the most energetic of the four calorimeter towers that make up the trigger tower and sums the energy in the EM sections of a 3×3 array of calorimeter towers around it. It checks the longitudinal shower shape by applying cuts on the fraction of the energy in the different EM layers. The transverse shower shape is characterized by the energy deposition pattern in the third EM layer. The difference between the energies in concentric regions covering 0.25×0.25 and 0.15×0.15 in $\eta \times \phi$ must be consistent with an electron. Level 2 also imposes an isolation condition requiring

$$\frac{\sum_i E_i \sin \phi_i - p_T}{p_T} < 0.15, \quad (10)$$

where the sum runs over all cells within a cone of radius $R = \sqrt{\Delta \phi^2 + \Delta \eta^2} = 0.4$ around the electron direction and p_T is the transverse momentum of the electron [27].

The p_T of the electron computed at level 2 is based on its energy and the z position of the interaction vertex measured by the level 0 hodoscopes. Level 2 accepts events that have a minimum number of EM clusters that satisfy the shape cuts and have p_T above a preprogrammed threshold. Figure 12




FIG. 13. Efficiency of a 15 GeV level 2 p_T requirement. The arrow indicates the cut applied in the final event selection.

shows the measured relative efficiency of the level 2 electron filter versus electron p_T for a level 2 p_T threshold of 20 GeV. We determine this efficiency using Z data taken with a lower threshold value (16 GeV). The efficiency is the fraction of electrons above a level 2 p_T threshold of 20 GeV. The curve is the parametrization used in the fast Monte Carlo simulation.

Level 2 also computes the missing transverse momentum based on the energy registered in each calorimeter cell and the vertex z position. We determine the efficiency curve for a 15 GeV level 2 p_T requirement from data taken without the level 2 p_T condition. Figure 13 shows the measured efficiency versus $p_T(\nu)$. The curve is the parametrization used in the fast Monte Carlo simulation.

B. Reconstruction

1. Electron

We identify electrons as clusters of adjacent calorimeter cells with significant energy deposits. Only clusters with at least 90% of their energy in the EM section and at least 60% of their energy in the most energetic calorimeter tower are considered as electron candidates. For most electrons we also reconstruct a track in the CDC or FDC that points towards the centroid of the cluster.

We compute the electron energy $E(e)$ from the signals in all cells of the EM layers and the first FH layer in a window covering 0.5×0.5 in $\eta \times \phi$ and centered on the tower which registered the highest fraction of the electron energy. In the computation we use the sampling weights and calibration constants determined using the test beam data (Sec. III B 3) except for the offset δ_{EM} , which we take from an in situ calibration (Sec. VI D), i.e., $\delta_{EM} = -0.16$ GeV for electrons in the CC.

The calorimeter shower centroid position $(x_{cal}, y_{cal}, z_{cal})$, the center of gravity of the track $(x_{trk}, y_{trk}, z_{trk})$, and the proton beam trajectory define the electron direction. The shower centroid algorithm is documented in Appendix B. The center of gravity of the CDC track is defined by the mean hit coordinates of all the delay line hits on the track.

The calibration of the measured z coordinates contributes a significant systematic uncertainty to the W boson mass measurement and is described in Appendixes A and B. Using tracks from many events reconstructed in the vertex drift chamber, we measure the beam trajectory for every run. The closest approach to the beam trajectory of the line through shower centroid and track center of gravity defines the position of the interaction vertex $(x_{\text{vtx}}, y_{\text{vtx}}, z_{\text{vtx}})$. In $Z \rightarrow ee$ events we may have two electron candidates with tracks. In this case we take the point midway between the vertex positions determined from each electron as the interaction vertex. Using only the electron track to determine the position of the interaction vertex, rather than all tracks in the event, makes the resolution of this measurement less sensitive to the luminosity and avoids confusion between vertices in events with more than one $p\bar{p}$ interaction.

We then define the azimuth $\phi(e)$ and the colatitude $\theta(e)$ of the electron using the vertex and the shower centroid positions:

$$\tan \phi(e) = \frac{y_{\text{cal}} - y_{\text{vtx}}}{x_{\text{cal}} - x_{\text{vtx}}}, \quad (11)$$

$$\tan \theta(e) = \frac{\sqrt{x_{\text{cal}}^2 + y_{\text{cal}}^2} - \sqrt{x_{\text{vtx}}^2 + y_{\text{vtx}}^2}}{z_{\text{cal}} - z_{\text{vtx}}}. \quad (12)$$

Neglecting the electron mass, the momentum of the electron is given by

$$\vec{p}(e) = E(e) \begin{pmatrix} \sin \theta(e) \cos \phi(e) \\ \sin \theta(e) \sin \phi(e) \\ \cos \theta(e) \end{pmatrix}. \quad (13)$$

2. Recoil

We reconstruct the transverse momentum of all particles recoiling against the W or Z boson by taking the vector sum

$$\vec{u}_T = \sum_i E_i \sin \theta_i \begin{pmatrix} \cos \phi_i \\ \sin \phi_i \end{pmatrix}, \quad (14)$$

where the sum runs over all calorimeter cells that were read out, except those that belong to electron clusters. E_i are the cell energies, and ϕ_i and θ_i are the azimuth and colatitude of the center of cell i with respect to the interaction vertex.

3. Derived quantities

In the case of $Z \rightarrow ee$ decays, we define the dielectron momentum

$$\vec{p}(ee) = \vec{p}(e_1) + \vec{p}(e_2) \quad (15)$$

and the dielectron invariant mass

$$m(ee) = \sqrt{2E(e_1)E(e_2)(1 - \cos \omega)}, \quad (16)$$

where ω is the opening angle between the two electrons. It is useful to define a coordinate system in the plane transverse to the beam that depends only on the electron directions. We




FIG. 14. Illustration of momentum vectors in the transverse plane for $Z \rightarrow ee$ candidates. The vectors drawn with thick lines are directly measured.

follow the conventions first introduced by UA2 [10] and call the axis along the inner bisector of the two electrons the η axis and the axis perpendicular to that the ξ axis. Projections on these axes are denoted with subscripts η or ξ . Figure 14 illustrates these definitions.

In case of $W \rightarrow e\nu$ decays we define the transverse neutrino momentum

$$\vec{p}_T(\nu) = -\vec{p}_T(e) - \vec{u}_T \quad (17)$$

and the transverse mass [Eq. (6)]. Useful quantities are the projection of the transverse recoil momentum on the electron direction,

$$u_{\parallel} = \vec{u}_T \cdot \hat{p}_T(e), \quad (18)$$

and the projection on the direction perpendicular to the electron direction,

$$u_{\perp} = \vec{u}_T \cdot [\hat{p}_T(e) \times \hat{z}]. \quad (19)$$

Figure 15 illustrates these definitions.

C. Electron identification

1. Fiducial cuts

To ensure a uniform response we accept only electron candidates that are well separated in the azimuth ($\Delta\phi$) from the calorimeter module boundaries in the CC-EM section




FIG. 15. Illustration of momentum vectors in the transverse plane for $W \rightarrow e\nu$ candidates. The vectors drawn with thick lines are directly measured.

and from the edges of the calorimeter by cutting on $\Delta\phi$ and z_{cal} . We also remove electrons for which the z position of the track center of gravity is near the edge of the CDC. For electrons in the EC-EM section we cut on the index of the most energetic tower, i_η . Tower 15 covers $1.4 < \eta < 1.5$ with respect to the detector center, and tower 25 covers $2.4 < \eta < 2.5$.

2. Quality variables

We test how well the shape of a cluster agrees with that expected for an electromagnetic shower by computing a quality variable (χ^2) for all cell energies using a 41-dimensional covariance matrix. The covariance matrix was determined from GEANT [28] based simulations [29].

To determine how well a track matches a cluster, we extrapolate the track to the third EM layer in the calorimeter and compute the distance between the extrapolated track and the cluster centroid in the azimuthal direction, Δs , and in the z direction, Δz . The variable

$$\sigma_{\text{trk}}^2 = \left(\frac{\Delta s}{\delta s} \right)^2 + \left(\frac{\Delta z}{\delta z} \right)^2 \quad (20)$$

quantifies the quality of the match. In the EC-EM section z is replaced by r , the radial distance from the center of the detector. The parameters $\delta s = 0.25$ cm, $\delta z = 2.1$ cm, and $\delta r = 1.0$ cm are the resolutions with which Δs , Δz , and Δr are measured, as determined with the electrons from $W \rightarrow e\nu$ decays.

In the EC, electrons must have a matched track in the forward drift chamber. In the CC, we define ‘‘tight’’ and ‘‘loose’’ criteria. The tight criteria require a matched track in the CDC. The loose criteria do not require a matched track and help increase the electron finding efficiency for $Z \rightarrow ee$ decays.

The isolation fraction is defined as

$$f_{\text{iso}} = \frac{E_{\text{cone}} - E_{\text{core}}}{E_{\text{core}}}, \quad (21)$$

where E_{cone} is the energy in a cone of radius $R = \sqrt{\Delta\phi^2 + \Delta\eta^2} = 0.4$ around the direction of the electron, summed over the entire depth of the calorimeter and E_{core} is the energy in a cone of $R = 0.2$, summed over the EM calorimeter only.

Figure 16 shows the distributions of the three quality variables for electrons in the CC with the arrow showing the cut values. Table I summarizes the electron selection criteria.

D. Data samples

The data were taken during the 1994–1995 Tevatron run. After the removal of runs in which parts of the detector were not operating adequately, they amount to an integrated luminosity of about 82 pb^{-1} . We select W decay candidates by requiring the following.

- level 1: $p\bar{p}$ interaction
- Main Ring veto
- EM trigger tower above 10 GeV




FIG. 16. Distributions of the electron identification variables. The arrows indicate the cut values.

- level 1.5: ≥ 1 EM cluster above 15 GeV
- level 2: electron candidate with $p_T > 20$ GeV
momentum imbalance $p_T > 15$ GeV
- off line: ≥ 1 tight electron candidate in the CC
 $p_T(e) > 25$ GeV
 $p_T(\nu) > 25$ GeV
 $u_T < 15$ GeV

We select Z decay candidates by requiring the following.

- level 1: $p\bar{p}$ interaction
 ≥ 2 EM trigger towers above 7 GeV
- level 1.5: ≥ 1 EM cluster above 10 GeV
- level 2: ≥ 2 electron candidates with $p_T > 20$ GeV
- off line: ≥ 2 electron candidates
 $p_T(e) > 25$ GeV
 $70 < m(ee) < 110$ GeV

We accept $Z \rightarrow ee$ decays with at least one electron candidate in the CC and the other in the CC or EC. One CC candidate must pass the tight electron selection criteria. If the other candidate is also in the CC, it may pass only the loose criteria. We use the 2179 events with both electrons in the CC (CC/CC Z sample) to calibrate the calorimeter response to electrons (Sec. VI). These events need not pass the Main Ring veto cut because Main Ring background does not affect the EM calorimeter. The 2341 events for which both electrons have tracks and which pass the Main Ring veto (CC/CC+EC Z sample) serve to calibrate the recoil momentum response (Sec. VII). Table II summarizes the data samples.

TABLE I. Electron selection criteria.

Variable	CC (loose)	CC (tight)	EC (tight)
Fiducial cuts	$ \Delta\phi > 0.02$ $ z_{\text{cal}} < 108$ cm	$ \Delta\phi > 0.02$ $ z_{\text{cal}} < 108$ cm $ z_{\text{trk}} < 80$ cm	$15 \leq i_\eta \leq 25$
Shower shape	$\chi^2 < 100$	$\chi^2 < 100$	$\chi^2 < 100$
Isolation	$f_{\text{iso}} < 0.15$	$f_{\text{iso}} < 0.15$	$f_{\text{iso}} < 0.15$
Track match		$\sigma_{\text{trk}} < 5$	$\sigma_{\text{trk}} < 10$

TABLE II. Number of W and Z candidate events.

Channel	Z→ee			W→eν
	CC/CC	CC/CC	CC/EC	CC
Fiducial region of electrons	CC/CC	CC/CC	CC/EC	CC
e quality (t=tight, l=loose)	t/l	t/t	t/t	t
Pass Main Ring veto	537	1225	1116	28 323
Fail Main Ring veto	107	310	268	

Figure 17 shows the luminosity of the colliding beams during the W and Z data collection.

On several occasions we use a sample of 295 000 random $p\bar{p}$ interaction events for calibration purposes. We collected these data concurrently with the W and Z signal data, requiring only a $p\bar{p}$ interaction at level 1. We refer to these data as “minimum bias events.”

V. FAST MONTE CARLO MODEL

A. Overview

The fast Monte Carlo model consists of three parts. First we simulate the production of the W or Z boson by generating the boson four-momentum and other characteristics of the event like the z position of the interaction vertex and the luminosity. The event luminosity is required for luminosity-dependent parametrizations in the detector simulation. Then we simulate the decay of the boson. At this point we know the true p_T of the boson and the momenta of its decay products. We then apply a parametrized detector model to these momenta in order to simulate the observed transverse recoil momentum and the observed electron momenta.

B. Vector boson production

In order to specify completely the production dynamics of vector bosons in $p\bar{p}$ collisions, we need to know the differential production cross section in mass Q , rapidity y , and transverse momentum q_T of the produced W bosons. To speed up the event generation, we factorize this into




FIG. 17. Luminosity distribution of the W (solid line) and the Z (●) samples.

$$\frac{d^3\sigma}{dq_T^2 dy dQ} \approx \frac{d^2\sigma}{dq_T^2 dy} \Big|_{Q^2=M_W^2} \times \frac{d\sigma}{dQ} \quad (22)$$

to generate q_T , y , and Q of the bosons.

For $p\bar{p}$ collisions, the vector boson production cross section is given by the parton cross section $\tilde{\sigma}_{i,j}$ convoluted with the parton distribution functions $f(x, Q^2)$ and summed over parton flavors i, j :

$$\frac{d^2\sigma}{dq_T^2 dy} = \sum_{i,j} \int dx_1 \int dx_2 f_i(x_1, Q^2) f_j(x_2, Q^2) \times \delta(sx_1x_2 - Q^2) \frac{d^2\tilde{\sigma}_{i,j}}{dq_T^2 dy}. \quad (23)$$

Several authors [30,31] have computed $d^2\sigma/dq_T^2 dy|_{Q^2=M_W^2}$ using a perturbative calculation [32] for the high- q_T regime and the Collins-Soper resummation formalism [33,34] for the low- q_T regime. We use the code provided by the authors of Ref. [30] and the Martin-Roberts-Sterling set A' (MRSA') parton distribution functions [35] to compute the cross section. We evaluate Eq. (23) separately for interactions involving at least one valence quark and for interactions involving two sea quarks.

The parton cross section is given by

$$\frac{d^2\tilde{\sigma}}{dq_T^2 dy} = \frac{\tilde{\sigma}_0}{4\pi\hat{s}} \left\{ \int d^2b e^{iq_T \cdot \vec{b}} \cdot \tilde{W}(b) \times e^{-S+Y} \right\}, \quad (24)$$

where $\tilde{\sigma}_0$ is the tree-level cross section, \hat{s} is the parton center-of-mass energy, and b is the impact parameter in transverse momentum space. \tilde{W} and Y are perturbative terms, and S parametrizes the nonperturbative physics. In the notation of Ref. [30],

$$S = \left[g_1 + g_2 \ln\left(\frac{Q}{2Q_0}\right) \right] b^2 + g_1 g_3 \ln(100x_1x_2)b, \quad (25)$$

where Q_0 is a cutoff parameter and x_1 and x_2 are the momentum fractions of the initial state partons. The parameters g_1 , g_2 , and g_3 have to be determined experimentally (Sec. VIII).

We use a Breit-Wigner curve with mass-dependent width for the line shape of the W boson. The intrinsic width of the W is $\Gamma_W = 2.062 \pm 0.059$ GeV [36]. The line shape is skewed due to the momentum distribution of the quarks inside the proton and antiproton. The mass spectrum is given by

$$\frac{d\sigma}{dQ} = \mathcal{L}_{q\bar{q}}(Q) \frac{Q^2}{(Q^2 - M_W^2)^2 + Q^4 \Gamma_W^2 / M_W^2}. \quad (26)$$

We call

$$\mathcal{L}_{q\bar{q}}(Q) = \frac{2Q}{s} \sum_{i,j} \int_{Q^2/s}^1 \frac{dx}{x} f_i(x, Q^2) f_j(Q^2/sx, Q^2) \quad (27)$$

TABLE III. Parton luminosity slope β and fraction of sea-sea interactions f_{ss} in the W and Z production model. The β value is given for $W \rightarrow e\nu$ decays with the electron in the CC and for $Z \rightarrow ee$ decays with both electrons in the CC.

	W production		f_{ss}
	Z production β (GeV^{-1})	β (GeV^{-1})	
MRSA' [35]	3.6×10^{-3}	8.6×10^{-3}	0.207
CTEQ3M [42]	3.3×10^{-3}	8.7×10^{-3}	0.203
CTEQ2M [43]		8.8×10^{-3}	0.203
MRSD-' [44]	3.8×10^{-3}	9.6×10^{-3}	0.201

the parton luminosity. To evaluate it we generate $W \rightarrow e\nu$ events using the HERWIG Monte Carlo event generator [37], interfaced with PDFLIB [38], and select the events subject to the same kinematic and fiducial cuts as for the W and Z samples with all electrons in the CC. We plot the mass spectrum divided by the intrinsic line shape of the W boson. The result is proportional to the parton luminosity, and we parametrize the spectrum with the function [12]

$$\mathcal{L}_{q\bar{q}}(Q) = \frac{e^{-\beta Q}}{Q}. \quad (28)$$

Table III shows β for W and Z events for some modern parton distribution functions. The value of β depends on the rapidity distribution of the W bosons, which is restricted by the kinematic and fiducial cuts that we impose on the decay leptons. The values of β given in Table III are for the rapidity distributions of W and Z bosons that satisfy the kinematic and fiducial cuts given in Sec. IV. The uncertainty in β is about 0.001, due to Monte Carlo statistics and uncertainties in the acceptance.

To generate the boson four-momenta, we treat $d\sigma/dQ$ and $d^2\sigma/dq_T^2 dy$ as probability density functions and pick Q from the former and a pair of y and q_T values from the latter. For a fraction f_{ss} we use $d^2\sigma/dq_T^2 dy$ for interactions between two sea quarks. Their helicity is $+1$ or -1 with equal probability. For the remaining W bosons we use $d^2\sigma/dq_T^2 dy$ for interactions involving at least one valence quark. They always have helicity -1 . Finally, we pick the z position of the interaction vertex from a Gaussian distribution centered at $z=0$ with a standard deviation of 25 cm and a luminosity for each event from the histogram in Fig. 17.

C. Vector boson decay

At lowest order the W boson is fully polarized along the beam direction due to the V - A coupling of the charged current. The resulting angular distribution of the charged lepton in the W rest frame is given by

$$\frac{d\sigma}{d \cos \theta^*} \propto (1 - \lambda q \cos \theta^*)^2, \quad (29)$$

where λ is the helicity of the W with respect to the proton direction, q is the charge of the lepton, and θ^* is the angle between the charged lepton and proton beam directions in




FIG. 18. Polarization of the W produced in $p\bar{p}$ collisions if the quark comes from the proton (left) and if the antiquark comes from the proton (right). The thick arrows indicate the orientation of the particle spins.

the W rest frame. The spin of the W points along the direction of the incoming antiquark. Most of the time the quark comes from the proton and the antiquark from the antiproton, so that $\lambda = -1$. Only if both quark and antiquark come from the sea of the proton and antiproton is there a 50% chance that the quark comes from the antiproton and the antiquark from the proton and in that case $\lambda = 1$ (Fig. 18). We determine the fraction of sea-sea interactions, f_{ss} , using the parametrizations of the parton distribution functions given in PDFLIB [38].

When $\mathcal{O}(\alpha_s)$ processes are included, the boson acquires finite p_T and Eq. (29) is changed to [39]

$$\frac{d\sigma}{d \cos \theta_{CS}} \propto [1 + \alpha_1(q_T) \cos \theta_{CS} + \alpha_2(q_T) \cos^2 \theta_{CS}] \quad (30)$$

for W^+ bosons with $\lambda = -1$ and after integration over ϕ . The angle θ_{CS} in Eq. (30) is now defined in the Collins-Soper frame [40]. The values of α_1 and α_2 as a function of transverse boson momentum have been calculated at $\mathcal{O}(\alpha_s^2)$ [39] and are shown in Fig. 19. We have implemented the angular distribution given in Eq. (30) in the fast Monte Carlo model. The effect is smaller if the W bosons are selected with $u_T < 15$ GeV rather than for $u_T < 30$ GeV. The angular distribution of the leptons from $Z \rightarrow ee$ decays is also generated according to Eq. (30), but with α_1 and α_2 computed for $Z \rightarrow ee$ decays [39].

To check whether neglecting the correlations between the mass and the other parameters in Eq. (22) introduces an uncertainty, we use the HERWIG program to generate $W \rightarrow e\nu$ decays including the correlations neglected in our model. We




FIG. 19. Calculations of α_1 (solid line) and α_2 (dashed line) as a function of the transverse momentum of the W boson.




FIG. 20. Distribution of $\Delta R(e\gamma)$ of photons from $W \rightarrow e\nu\gamma$ decays that are reconstructed as separate objects (shaded area) and those that are not, either because they are too close to the electron or too low in energy (solid line).

determine all the input distributions (W boson line shape, differential cross section in p_T and y , and parton luminosity) from these events. We then feed these into our simulation. This generates events with the dynamics as in HERWIG, but without correlations between these input distributions. Alternatively, we directly feed the events generated by HERWIG into our simulation and only apply the parametrized detector model. The results from fits to the transverse mass spectra of these two samples differ by less than 15 ± 25 MeV. The uncertainty is due to the finite number of Monte Carlo events.

Radiation from the decay electron or the W boson biases the mass measurement. If the decay electron radiates a photon and the photon is well enough separated from the electron so that its energy is not included in the electron energy or if an on-shell W boson radiates a photon and therefore is off shell when it decays, the measured mass is biased low. We use the calculation of Ref. [41] to generate $W \rightarrow e\nu\gamma$ decays. The calculation gives the fraction of events in which a photon with energy $E(\gamma) > E_0$ is radiated, and the angular distribution and energy spectrum of the photons. Only radiation from the decay electron and the W boson, if the final state W is off shell, is included to order α . Radiation by the initial quarks or the W , if the final W is on shell, does not affect the mass of the $e\nu$ pair from the W decay. We use a minimum photon energy $E_0 = 50$ MeV, which means that in 30.6% of all W decays a photon with $E(\gamma) > 50$ MeV is radiated. Most of these photons are emitted close to the electron direction and cannot be separated from the electron in the calorimeter. For $Z \rightarrow ee$ decays there is a 66% probability that any one of the electrons radiates a photon with $E(\gamma) > 50$ MeV.

The separation of the electron and photon in the laboratory frame is

$$\Delta R(e\gamma) = \sqrt{[\phi(e) - \phi(\gamma)]^2 + [\eta(e) - \eta(\gamma)]^2}. \quad (31)$$

Figure 20 shows the calculated distribution of photons as a

function of $\Delta R(e\gamma)$. The shaded histogram in the figure shows the photons that are reconstructed as separate objects. If the photon and electron are close together, they cannot be separated in the calorimeter. The momentum of a photon with $\Delta R(e\gamma) < R_0$ is therefore added to the electron momentum, while for $\Delta R(e\gamma) \geq R_0$ a photon is considered separated from the electron and its momentum is added to the recoil momentum. We use $R_0 = 0.3$, which is the approximate size of the window in which the electron energy is measured. This procedure has been verified to give the same results as an explicit GEANT simulation of radiative W decays. In only about 3.5% of the $W \rightarrow e\nu$ decays does the photon separate far enough from the electron, i.e., $\Delta R(e\gamma) > R_0$, to cause a mismeasurement of the transverse mass.

W boson decays through the channel $W \rightarrow \tau\nu \rightarrow e\nu\bar{\nu}$ are topologically indistinguishable from $W \rightarrow e\nu$ decays. We therefore include these decays in the W decay model, properly accounting for the polarization of the τ leptons in the decay angular distributions. The fraction of W bosons that decay in this way is $B(\tau \rightarrow e\nu\bar{\nu}) / [1 + B(\tau \rightarrow e\nu\bar{\nu})] = 0.151$.

We let the generated W bosons decay with an angular distribution corresponding to their helicity. For 15.1% of the W bosons, the decay is to $\tau\nu \rightarrow e\nu\bar{\nu}$. For 30.6% of the remaining W bosons, a photon is radiated. For 66% of the Z bosons, the decay is to $e^+e^-\gamma$ and for the remainder to e^+e^- .

D. Detector model

The detector simulation uses a parametrized model for response and resolution to obtain a prediction for the distribution of the observed electron and recoil momenta.

When simulating the detector response to an electron of energy E_0 , we compute the observed electron energy as

$$E(e) = \alpha_{EM}E_0 + \Delta E(\mathcal{L}, u_{\parallel}) + \sigma_{EM} \cdot X, \quad (32)$$

where α_{EM} is the response of the electromagnetic calorimeter, ΔE is the energy due to particles from the underlying event within the electron window (parametrized as a function of luminosity \mathcal{L} and u_{\parallel}), σ_{EM} is the energy resolution of the electromagnetic calorimeter, and X is a random variable from a normal parent distribution with zero mean and unit width.

The transverse energy measurement depends on the measurement of the electron direction as well. We determine the shower centroid position by intersecting the line defined by the event vertex and the electron direction with a cylinder coaxial with the beam and 91.6 cm in radius (the radial center of the EM3 layer). We then smear the azimuthal and z coordinate of the intersection point by their resolutions. We determine the z coordinate of the center of gravity of the CDC track by intersecting the same line with a cylinder of 62 cm radius, the mean radial position of all delay lines in the CDC, and smearing by the resolution. The measured angles are then obtained from the smeared points as described in Sec. IV B 1.

The model for the particles recoiling against the W has two components: a ‘‘hard’’ component that models the p_T of the W and a ‘‘soft’’ component that models detector noise




FIG. 21. Distribution of $z_{\text{vtx}}(e_1) - z_{\text{vtx}}(e_2)$ for the $Z \rightarrow ee$ sample (●) and the fast Monte Carlo simulation (solid line).

and pileup. Pileup refers to the effects of additional $p\bar{p}$ interactions in the same or previous beam crossings. For the soft component we use the transverse momentum balance \vec{p}_T from a minimum bias event recorded in the detector. The observed recoil p_T is then given by

$$\vec{u}_T = -(R_{\text{rec}} q_T + \sigma_{\text{rec}} \cdot X) \hat{q}_T - \Delta u_{\parallel}(\mathcal{L}, u_{\parallel}) \hat{p}_T(e) + \alpha_{\text{mb}} \vec{p}_T, \quad (33)$$

where q_T is the generated value of the boson transverse momentum, R_{rec} is the (in general momentum-dependent) response, σ_{rec} is the resolution of the calorimeter, Δu_{\parallel} is the transverse energy flow into the electron window (parametrized as a function of luminosity and u_{\parallel}), and α_{mb} is a correction factor that allows us to adjust the resolution to the data. The quantity Δu_{\parallel} is different from the energy added to the electron, ΔE , because of the zero suppression in the calorimeter readout.

We simulate selection biases due to the trigger requirements and the electron isolation by accepting events with the estimated efficiencies. Finally, we compute all the derived quantities from these observables and apply fiducial and kinematic cuts.

VI. ELECTRON MEASUREMENT

A. Angular resolutions

The resolution for the z coordinate of the track center of gravity, z_{trk} , is determined from the $Z \rightarrow ee$ sample. Both electrons originate from the same interaction vertex, and therefore the difference between the interaction vertices reconstructed from the two electrons separately, $z_{\text{vtx}}(e_1) - z_{\text{vtx}}(e_2)$, is a measure of the resolution with which the electrons point back to the vertex. The points in Fig. 21 show the distribution of $z_{\text{vtx}}(e_1) - z_{\text{vtx}}(e_2)$ observed in the CC/CC Z sample with tracks required for both electrons.

A Monte Carlo study based on single electrons generated with a GEANT simulation shows that the resolution of the shower centroid algorithm can be parametrized as

$$\sigma(z_{\text{cal}}) = [a + b\lambda(e)] + [c + d\lambda(e)]z_{\text{cal}}, \quad (34)$$




FIG. 22. Transverse energy flow into 1×5 tower segments as a function of azimuthal separation from the electron in the W sample.

where $\lambda(e) = |\theta(e) - 90^\circ|$, $a = 0.33$ cm, $b = 5.2 \times 10^{-3}$ cm, $c = 4.2 \times 10^{-4}$, and $d = 7.5 \times 10^{-5}$. We then tune the resolution function for z_{trk} in the fast Monte Carlo simulation so that it reproduces the shape of the $z_{\text{vtx}}(e_1) - z_{\text{vtx}}(e_2)$ distribution observed in the data. We find that a resolution function consisting of two Gaussians 0.31 and 1.56 cm wide, with 6% of the area under the wider Gaussian, fits the data well. The histogram in Fig. 21 shows the Monte Carlo prediction for the best fit, normalized to the same number of events as the data. The W mass measurement is very insensitive to these resolutions. The uncertainties in the resolution parameters cause less than 5 MeV uncertainty in the fitted W mass.

The calibration of the z -position measurements from the CDC and calorimeter is described in Appendix A. We quantify the calibration uncertainty in terms of scale factors $\alpha_{\text{CDC}} = 0.988 \pm 0.001$ and $\alpha_{\text{CC}} = 0.9980 \pm 0.0005$ for the z coordinate. The uncertainties in these scale factors lead to a finite uncertainty in the W mass measurement.

B. Underlying event energy

The energy in an array of 5×5 towers in the four EM layers and the first FH layer around the most energetic tower of an electron cluster is assigned to the electron. This array contains the entire energy deposited by the electron shower plus some energy from other particles. The energy in the window is excluded from the computation of \vec{u}_T . This causes a bias in u_{\parallel} , the component of \vec{u}_T along the direction of the electron. For $p_T(W) \ll M_W$,

$$m_T \approx 2p_T(e) + u_{\parallel}, \quad (35)$$

so that this bias propagates directly into a bias in the transverse mass. We call this bias Δu_{\parallel} . It is equal to the momentum flow observed in the EM and first FH sections of a 5×5 array of calorimeter towers.



We use the W and Z data samples to measure Δu_{\parallel} . For every electron in the W and Z samples, we compute the energy flow into an azimuthal ring of calorimeter towers, five towers wide in η and centered on the tower with the largest

TABLE IV. Δu_{\parallel} for the W and Z event samples.

Event sample	$\Sigma E_{1 \times 5} / \cosh \eta(e)$ (MeV)	Δu_{\parallel} (MeV)
W	95.8 ± 0.4	$479 \pm 2 \pm 6$
Z	93.6 ± 1.3	$468 \pm 7 \pm 6$

fraction of the electron energy. For every electron we plot the transverse energy flow into one-tower-wide azimuthal segments of this ring as a function of the azimuthal separation $|\Delta\phi|$ between the center of the segment and the electron shower centroid. The energy flow $\Sigma E_{1 \times 5}$ is computed as the sum of all energy deposits in the four EM layers and the first FH layer in the 1×5 tower segment. Figure 22 shows the transverse energy flow $\Sigma E_{1 \times 5} / \cosh \eta(e)$ versus $|\Delta\phi|$ for the electrons in the W sample with $u_T < 15$ GeV. For small $|\Delta\phi|$ we see a substantial energy flow from the electron shower and for larger $|\Delta\phi|$ a constant noise level. The electron shower is contained in a window of $|\Delta\phi| < 0.2$. We estimate the energy flow into the 5×5 tower window around the electron from the energy flow into segments of the azimuthal ring with $|\Delta\phi| > 0.2$. The level of energy flow is sensitive to the isolation cut. The region $0.2 < |\Delta\phi| < 0.4$, which is used for the isolation variable, is maximally biased by the cut; the region $0.4 < |\Delta\phi| < 0.6$, which is close to the electron but outside the isolation region, is minimally biased. We expect the energy flow under the electron to lie somewhere in between the energy flow into these two regions. We therefore compute Δu_{\parallel} based on the average transverse energy flow into both regions and assign a systematic error equal to half the difference between the two regions. We repeat the same analysis for the electrons in the CC/CC Z sample. The results are tabulated in Table IV. We find $\Delta u_{\parallel} = 479 \pm 2(\text{stat}) \pm 6(\text{syst})$ MeV for W events with $u_T < 15$ GeV. For the Z sample, Δu_{\parallel} is 11 ± 7 MeV lower. Figure 23 shows the spectrum of Δu_{\parallel} .

At higher luminosity the average number of interactions per event increases and therefore Δu_{\parallel} increases. This is shown in Fig. 24. The mean value of Δu_{\parallel} increases by 11.2 MeV per $10^{30} \text{ cm}^{-2} \text{ s}^{-1}$. The underlying event energy flow

FIG. 23. Distribution of Δu_{\parallel} in the W signal sample.FIG. 24. The luminosity dependence of $\langle \Delta u_{\parallel} \rangle$.

into the electron window also depends on u_{\parallel} . Figure 25 shows $\langle \Delta u_{\parallel}(0, u_{\parallel}) \rangle$, the mean value for Δu_{\parallel} corrected back to zero luminosity, as a function of u_{\parallel} . In the fast Monte Carlo model a value Δu_{\parallel} is picked from the distribution shown in Fig. 23 for every event and then corrected for u_{\parallel} and luminosity dependences.


The measured electron energy is biased upwards by the additional energy ΔE in the window from the underlying event. ΔE is not equal to Δu_{\parallel} because the additional energy deposited by the electron may lift some cells that would have been zero suppressed in the calorimeter readout above the zero-suppression threshold. Therefore

$$\Delta E = \Delta u_{\parallel} - \Delta_{\text{ped}}, \quad (36)$$

where $\Delta_{\text{ped}} = 212 \pm 25$ MeV is a correction for the pedestal shift introduced by the zero suppression in the calorimeter readout. This is determined by superimposing single electrons simulated with a GEANT simulation on minimum bias events that were recorded without zero suppression in the calorimeter readout. Most of this bias cancels in the W to Z mass ratio so that the W mass measurement is not sensitive to Δ_{ped} .

C. u_{\parallel} efficiency

The efficiency for electron identification depends on their environment. Well-isolated electrons are identified correctly more often than electrons near other particles. Therefore W decays in which the electron is emitted in the same direction as the particles recoiling against the W are selected less often than W decays in which the electron is emitted in the direction opposite the recoiling particles. This causes a bias in the lepton p_T distributions, shifting $p_T(e)$ to larger values and $p_T(\nu)$ to lower values, whereas the m_T distribution is only slightly affected.

FIG. 25. Variation of $\langle \Delta u_{\parallel} \rangle$ as a function of u_{\parallel} . The region between the arrows is populated by the W sample.




FIG. 26. Transverse energy flow into 1×5 tower segments as a function of the azimuthal separation from the electron for the electrons from $W \rightarrow e\nu$ decays (●) and the superimposed Monte Carlo electron sample (solid line).

We estimate the electron finding efficiency as a function of u_{\parallel} by superimposing Monte Carlo electrons, simulated using the GEANT program, onto the events from our W signal sample. We use the W sample in order to ensure that the underlying event is correctly modeled. The sample of superimposed electrons, which are spatially separated from the electron that is already in the event, matches the data well. It is important that the superimposed sample model the transverse shower shape and isolation well, because these are the dominant effects that cause the efficiency to vary with u_{\parallel} . Figure 26 shows the transverse shower profile of the superimposed electron sample and the electron sample from W decays. Figure 27 shows the distribution of the isolation for the two electron samples in five u_{\parallel} regions. Figure 28 compares the mean isolation versus u_{\parallel} for the two samples.

We then apply the shower shape and isolation cuts used to select the W signal sample and determine the fraction of the electrons in the superimposed samples that pass all require-




FIG. 27. Isolation spectrum for five different u_{\parallel} regions: $u_{\parallel} < -15$ GeV, $-15 < u_{\parallel} < -5$ GeV, $-5 < u_{\parallel} < 5$ GeV, $5 < u_{\parallel} < 15$ GeV, and $u_{\parallel} > 15$ GeV (from top to bottom), for the electrons from $W \rightarrow e\nu$ decays (●) and the superimposed electron sample (shaded area).




FIG. 28. Mean isolation versus u_{\parallel} for the W electron sample (○) and the superimposed Monte Carlo electron sample (●).

ments as a function of u_{\parallel} . This efficiency is shown in Fig. 29. The line is a fit to a function of the form

$$\varepsilon(u_{\parallel}) = \varepsilon_0 \begin{cases} 1 & \text{for } u_{\parallel} < u_0, \\ 1 - s(u_{\parallel} - u_0) & \text{otherwise.} \end{cases} \quad (37)$$

The parameter ε_0 is an overall efficiency which is inconsequential for the W mass measurement, u_0 is the value of u_{\parallel} at which the efficiency starts to decrease as a function of u_{\parallel} , and s is the rate of decrease. We obtain the best fit for $u_0 = 3.85 \pm 0.55$ GeV and $s = 0.013 \pm 0.001$ GeV⁻¹. These two values are strongly correlated. The errors account for the finite number of superimposed Monte Carlo electrons.

D. Electron energy response

Equation (7) relates the reconstructed electron energy to the recorded calorimeter signals. Since the values for the constants were determined in a different setup, we determine the offset δ_{EM} and a scale α_{EM} , which essentially modifies A , *in situ* with collider data for resonances that decay to electromagnetically showering particles: $\pi^0 \rightarrow \gamma\gamma$, $J/\psi \rightarrow ee$, and $Z \rightarrow ee$. We use π^0 and J/ψ signals from an




FIG. 29. Electron selection efficiency as a function of u_{\parallel} .




FIG. 30. Background-subtracted m_{sym} distribution. The superimposed curve shows the Monte Carlo simulation.

integrated luminosity of approximately 150 nb^{-1} , accumulated during dedicated runs with low- p_T thresholds for EM clusters in the trigger.

The fast Monte Carlo simulation predicts the reconstructed electron energy

$$E(e) = \alpha_{\text{EM}} E_0 = A \sum_{i=1}^5 s_i a_i - \delta_{\text{EM}}, \quad (38)$$

where E_0 is the generated electron energy. To determine δ_{EM} and α_{EM} , we compare the observed resonances and Monte Carlo predictions as a function of α_{EM} and δ_{EM} .

The photons from the decay of π^0 's with $p_T > 1 \text{ GeV}$ cannot be separated in the calorimeter. There is about a 10% probability for each photon to convert to an e^+e^- pair in the material in front of the CDC. If both photons convert, we can identify π^0 decays as EM clusters in the calorimeter with two doubly ionizing tracks in the CDC. We measure the π^0 energy $E(\pi^0)$ in the calorimeter and the opening angle ω between the two photons using the two tracks. This allows us to compute the ‘‘symmetric mass’’




FIG. 31. Dielectron invariant mass spectrum for the $J/\psi \rightarrow ee$ sample (histogram) and the background sample (\bullet). The smooth curve is a fit to the data.




FIG. 32. Dielectron mass spectrum for the background data sample to the CC/CC Z sample. The fit is an exponential.

$$m_{\text{sym}} = E(\pi^0) \sqrt{\frac{1 - \cos \omega}{2}}, \quad (39)$$

which is equal to the invariant mass if both photons have the same energy and is larger for asymmetric decays. Figure 30 shows the background-subtracted spectrum of m_{sym} for π^0 candidates in the CC-EM section superimposed with a Monte Carlo prediction of the line shape.

Figure 31 shows the invariant mass spectrum of dielectron pairs in the J/ψ mass region. The smooth curve is the fit to a Gaussian line shape above the background predicted using a sample of EM clusters without CDC tracks. After correcting by $-0.08 \pm 0.08 \text{ GeV}$ for the underlying event energy, we measure a mass of $3.03 \pm 0.04(\text{stat}) \pm 0.19(\text{syst}) \text{ GeV}$. A Monte Carlo simulation of $p\bar{p} \rightarrow b\bar{b} + X$, $b \rightarrow J/\psi + X$ tells us that we expect to observe a mass

$$m_{\text{obs}} = \alpha_{\text{EM}} m_{J/\psi} + 0.56 \delta_{\text{EM}}. \quad (40)$$

Together with our measurement of m_{obs} , this restricts the allowed parameter space for α_{EM} and δ_{EM} . The π^0 and J/ψ analyses are described in detail in Ref. [12]. Figure 34, below, shows the 68% confidence level contours in α_{EM} and δ_{EM} obtained from these data.

Fixing the observed Z boson mass to the measured value [Eq. (3)] correlates the values allowed for α_{EM} and δ_{EM} . For a given δ_{EM} we determine α_{EM} so that the position of the Z peak predicted by the fast Monte Carlo simulation agrees with the data. To determine the scale factor that best fits the data, we perform a maximum likelihood fit to the $m(ee)$ spectrum between 70 and 110 GeV. In the resolution function we allow for an exponential background shape whose slope is fixed to $-0.037 \pm 0.002 \text{ GeV}^{-1}$, the value obtained from a sample of events with two EM clusters that fail the electron quality cuts (Fig. 32). The background normalization is allowed to float in the fit. This is sufficient, together with the π^0 and J/ψ data, to determine both α_{EM} and δ_{EM} .

Without relying on the low-energy data at all, we can extract α_{EM} and δ_{EM} from the Z data alone. The electrons from Z decays are not monochromatic, and therefore we can




FIG. 33. Distribution of $m(ee)$ versus f_Z for the CC-CC $Z \rightarrow ee$ sample.

make use of their energy spread to constrain α_{EM} and δ_{EM} simultaneously. For $\delta_{EM} \ll E(e_1) + E(e_2)$ we can write

$$m(ee) = \alpha_{EM} M_Z + f_Z \delta_{EM}, \quad (41)$$

where $f_Z = [E(e_1) + E(e_2)](1 - \cos \omega) / m(ee)$ and ω is the opening angle between the two electrons. We plot $m(ee)$ versus f_Z (Fig. 33) and compare it with the Monte Carlo predictions for the allowed values of α_{EM} and δ_{EM} using a binned maximum likelihood fit.

Using the constraints on α_{EM} and δ_{EM} from the Z data alone, we obtain the contour labeled ‘‘Z’’ in Fig. 34 and $\delta_{EM} = 0.02 \pm 0.36$ GeV. The uncertainty in this measurement of δ_{EM} is dominated by the statistical uncertainty due to the finite size of the Z sample.

The combined constraint from all three resonances is shown by the thick contour in Fig. 34. The π^0 and J/ψ contours essentially fix δ_{EM} , independent of α_{EM} . The requirement that the Z peak position agree with the known Z boson mass correlates α_{EM} and δ_{EM} . The contours in Fig. 34




FIG. 34. Sixty-eight percent (68%) confidence level contours in α_{EM} and δ_{EM} from the J/ψ , π^0 , and Z data. The inset shows an expanded view of the region where the χ^2 is minimized.




FIG. 35. Dielectron mass spectrum from the CC-CC Z sample. The superimposed curve shows the maximum likelihood fit and the shaded region the fitted background.

reflect only statistical uncertainties. The uncertainty in the π^0 and J/ψ contours is dominated by systematic effects in the underlying event corrections and the deviation of the test beam data from the assumed response at low energies. The double arrow in Fig. 34 represents the systematic uncertainty in δ_{EM} . We determine

$$\delta_{EM} = -0.16^{+0.03}_{-0.21} \text{ GeV}. \quad (42)$$

Figure 35 shows the $m(ee)$ spectrum for the CC/CC Z sample and the Monte Carlo spectrum that best fits the data for $\delta_{EM} = -0.16$ GeV. The χ^2 for the best fit to the CC/CC $m(ee)$ spectrum is 33.5 for 39 degrees of freedom. For

$$\alpha_{EM} = 0.9533 \pm 0.0008, \quad (43)$$

the Z peak position is consistent with the known Z boson mass. The error reflects the statistical uncertainty and the uncertainty in the background normalization. The background slope has no measurable effect on the result.

If we split the CC/CC Z sample into events with two tight electrons and events with a tight and a loose electron and fit them separately using the value of α_{EM} given in Eq. (43), we obtain

$$M_Z = 91.206 \pm 0.086 \text{ GeV} \quad (\text{tight/tight sample}), \quad (44)$$

$$M_Z = 91.145 \pm 0.148 \text{ GeV} \quad (\text{tight/loose sample}). \quad (45)$$

Figures 36(a) and 36(b) show the corresponding spectra and fits.

E. Electron energy resolution

Equation (9) gives the functional form of the electron energy resolution. We take the intrinsic resolution of the calorimeter, which is given by the sampling term s_{EM} , from the test beam measurements. The noise term n_{EM} is represented by the width of the ΔE distribution (Fig. 23). We measure the constant term c_{EM} from the Z line shape of the data. We




FIG. 36. Dielectron mass spectra from (a) the tight/tight and (b) the tight/loose CC-CC Z samples. The curves show the fitted Monte Carlo spectra.

fit a Breit-Wigner line shape convoluted with a Gaussian, whose width characterizes the dielectron mass resolution, to the Z peak. Figure 37 shows the width $\sigma_{m(ee)}$ of the Gaussian fitted to the Z peak predicted by the fast Monte Carlo simulation as a function of c_{EM} . The horizontal lines indicate the width of the Gaussian fitted to the CC/CC Z sample and its uncertainties, 1.75 ± 0.08 GeV. We find that Monte Carlo simulation and data agree if $c_{EM} = 0.0115^{+0.0027}_{-0.0036}$, as indicated by the arrows in Fig. 37. The measured Z mass does not depend on c_{EM} .

VII. RECOIL MEASUREMENT

A. Recoil momentum response

The detector response and resolution for particles recoiling against a W boson should be the same as for particles recoiling against a Z boson. For $Z \rightarrow ee$ events, we can measure the transverse momentum of the Z from the e^+e^- pair, $p_T(ee)$, into which it decays and from the recoil momentum u_T in the same way as for $W \rightarrow e\nu$ events. By comparing $p_T(ee)$ and u_T we calibrate the recoil response relative to the electron response.

The recoil momentum is carried by many particles, mostly hadrons, with a wide momentum spectrum. Since the response of calorimeters to hadrons tends to be nonlinear and the recoil particles are distributed all over the calorimeter, including module boundaries with reduced response, we ex-




FIG. 37. Dielectron mass resolution versus the constant term c_{EM} .




FIG. 38. Recoil momentum response in the Monte Carlo Z sample as a function of q_T .

pect a momentum-dependent response function with values below unity. In order to fix the functional form of the recoil momentum response, we study the response predicted by a Monte Carlo $Z \rightarrow ee$ sample obtained using the HERWIG program and a GEANT-based detector simulation. We project the reconstructed transverse recoil momentum onto the direction of motion of the Z and define the response as

$$R_{\text{rec}} = \frac{|\vec{u}_T \cdot \hat{q}_T|}{q_T}, \quad (46)$$

where q_T is the generated transverse momentum of the Z boson. Figure 38 shows this response as a function of q_T . A response function of the form

$$R_{\text{rec}} = \alpha_{\text{rec}} + \beta_{\text{rec}} \log(q_T/\text{GeV}) \quad (47)$$

fits the response predicted by GEANT with $\alpha_{\text{rec}} = 0.713 \pm 0.006$ and $\beta_{\text{rec}} = 0.046 \pm 0.002$. This functional form also describes the jet energy response of the D0 calorimeter.




FIG. 39. Average $p_\eta(ee) + u_\eta$ versus $p_\eta(ee)$ for the Z data (●) and the fast Monte Carlo simulation (○).




FIG. 40. χ_0^2+1 contour for the recoil momentum response parameters.

To measure the recoil response from the collider data, we use the CC/CC+EC Z sample. We allow one of the leptons from the $Z \rightarrow ee$ decay to be in the CC or EC, so that the rapidity distribution of the Z bosons approximates that of the W bosons. We require both leptons to satisfy the tight electron criteria. This reduces the background for the topology with one lepton in the EC. We also require the Main Ring veto as for the W sample (Sec. IV).

We project the transverse momenta of the recoil, u_T , and the Z as measured by the two electrons, $p_T(ee)$, on the inner bisector of the electron directions (η axis), as shown in Fig. 14. By projecting the momenta on an axis that is independent of any energy measurement, noise contributions to the momenta average to zero and do not bias the result. We bin the data in $p_\eta(ee)$ and plot the mean of the sum of the two projections, $u_\eta + p_\eta(ee)$, versus the mean of $p_\eta(ee)$ (Fig. 39). We perform a two-dimensional χ^2 fit for the two parameters by comparing the data to predictions of the fast Monte Carlo model for different values of α_{rec} and β_{rec} . Figure 39 also shows the prediction of the Monte Carlo model for the values of the parameters that give the best fit. Figure 40 shows the contour for $\chi^2 = \chi_0^2 + 1$. The best fit ($\chi_0^2 = 5$ for eight degrees of freedom) is achieved for $\alpha_{\text{rec}} = 0.693 \pm 0.060$ and $\beta_{\text{rec}} = 0.040 \pm 0.021$. The two parameters are strongly correlated with a correlation coefficient $\rho = -0.979$.

B. Recoil momentum resolution

We parametrize the resolution for the hard component of the recoil as

$$\sigma_{\text{rec}} = s_{\text{rec}} \sqrt{u_T}, \quad (48)$$

where s_{rec} is a tunable parameter.

The soft component of the recoil is modeled by the transverse momentum balance \not{p}_T from minimum bias events, multiplied by a correction factor α_{mb} [Eq. (33)]. This automatically models the effects of detector resolution and pileup. To model the pileup correctly as a function of lumi-




FIG. 41. Width of the η -balance distribution versus $p_\eta(ee)$ for the Z data (\bullet) and the fast Monte Carlo simulation (\circ).

osity, we need to take the minimum bias events at the same luminosity as the W events. At a given luminosity the mean number of interactions in minimum bias events is always smaller than the mean number of interactions in W events. To model the detector resolution correctly, the minimum bias events must have the same interaction multiplicity spectrum as the W events. We therefore weight the minimum bias events so that their interaction multiplicity approximates that of the W events. As a measure of the interaction multiplicity on an event-by-event basis, we use the multiplicity of vertices reconstructed from the tracks in the CDC and the timing structure of the level 0 hodoscope signals [45].

We tune the two parameters s_{rec} and α_{mb} using the CC/CC+EC Z sample. The width of the spectrum of the η balance, $u_\eta/R_{\text{rec}} + p_\eta(ee)$, is a measure of the recoil momentum resolution. Figure 41 shows this width σ_η as a function of $p_\eta(ee)$. The contribution of the electron momentum resolution to the width of the η balance is negligibly small. The contribution of the recoil momentum resolution grows with $p_\eta(ee)$, while the contribution from the minimum bias




FIG. 42. χ_0^2+1 contour for the recoil resolution parameters α_{mb} and s_{rec} .




FIG. 43. η -balance distribution for the Z data (●) and the fast Monte Carlo simulation (solid line).

p_T is independent of $p_\eta(ee)$. This allows us to determine s_{rec} and α_{mb} simultaneously and without sensitivity to the electron resolution by comparing the width of the η balance predicted by the Monte Carlo model with that observed in the data in bins of $p_\eta(ee)$. We perform a χ^2 fit comparing Monte Carlo and collider data. Figure 42 shows contours of constant χ^2 in the α_{mb} - s_{rec} plane. The best agreement ($\chi_0^2 = 10.3$ for eight degrees of freedom) occurs for $s_{rec} = 0.49 \pm 0.14 \text{ GeV}^{1/2}$ and $\alpha_{mb} = 1.032 \pm 0.028$ with a correlation coefficient $\rho = -0.60$ for the two parameters. The ξ balance $u_\xi/R_{rec} + p_\xi(ee)$ is more sensitive to the electron momentum resolution and is affected by changes in s_{rec} and α_{mb} in the same way. We use it as a cross-check only.

Figure 43 shows the spectrum of $u_\eta/R_{rec} + p_\eta(ee)$ from the CC/CC+EC Z data sample and from the fast Monte Carlo model with the tuned recoil resolution and response parameters. Figure 44 shows the corresponding distributions for $u_\xi/R_{rec} + p_\xi(ee)$. In both cases the agreement between data and Monte Carlo simulation is good. A Kolmogorov-Smirnov test [46] gives confidence levels of $\kappa = 0.33$ and




FIG. 44. ξ -balance distribution for the Z data (●) and the fast Monte Carlo simulation (solid line).




FIG. 45. Transverse energy flow in the W (●) and Z (solid line) data.

0.37 that the Monte Carlo and data spectra derive from the same parent distribution. A χ^2 test gives $\chi^2 = 25$ and 37, respectively, for 40 bins.

Figure 45 shows the overall energy flow transverse to the beam direction measured by the sum $S_T = \sum_i E_i \sin \theta_i$ over all calorimeter cells except cells belonging to an electron cluster. For W events, $\langle S_T \rangle = 98.7 \pm 0.3 \text{ GeV}$, and for Z events, $\langle S_T \rangle = 91.0 \pm 0.9 \text{ GeV}$. Increased transverse energy flow leads to a worse recoil momentum resolution, and therefore we need to correct the value of α_{mb} for the W sample to account for this difference. Figure 46 relates transverse energy flow S_T to resolution σ_T for a minimum bias event sample. The resolution for measuring transverse momentum balance along any direction is

$$\sigma_T(S_T) = 1.42 \text{ GeV} + 0.15 \sqrt{S_T \text{ GeV}} + 0.007 S_T \quad (49)$$

for minimum bias events. The different energy flows in W and Z events lead to a correction to α_{mb} of $\sigma_T(98.7 \text{ GeV})/\sigma_T(91.0 \text{ GeV}) = 1.03 \pm 0.01$. The uncertainty




FIG. 46. Resolution for transverse momentum balance σ_T versus the transverse energy flow S_T for minimum bias events (●). The smooth curve is a fit [Eq. (49)].




FIG. 47. u_{\parallel} spectrum for the W data (●) and the Monte Carlo simulation (solid line).

reflects the uncertainties in the determination of $\langle S_T \rangle$. This uncertainty does not correlate with s_{rec} .

Z bosons are not intrinsically produced with less energy flow in the underlying event than W bosons. Rather, the requirement of two reconstructed isolated electrons biases the event selection in the Z sample towards events with lower-energy flow compared to the events in the W sample which have only one electron. We demonstrate this by loosening the electron identification requirements for one of the electrons in the Z sample. We use events that were collected using less restrictive trigger conditions for which at level 2 only one of the electron candidates must satisfy the shape and isolation requirements. We find that if all electron quality cuts are removed for one electron, S_T increases by 7%, consistent with the ratio of the S_T values in the W and Z samples.

C. Comparison with W data

We compare the recoil momentum distribution in the W data to the predictions of the fast Monte Carlo model, which




FIG. 48. u_{\perp} spectrum for the W data (●) and the Monte Carlo simulation (solid).




FIG. 49. Recoil momentum (u_T) spectrum for the W data (●) and the Monte Carlo simulation (solid line). The arrow shows the location of the cut.

includes the parameters determined in this section and Sec. VI. Figure 47 compares the u_{\parallel} spectra from Monte Carlo and W data. The mean u_{\parallel} for the W data is -0.64 ± 0.03 GeV, and for the Monte Carlo prediction including backgrounds it is -0.61 ± 0.01 GeV, in very good agreement. For the $W \rightarrow e\nu$ signal only, the Monte Carlo model predicts a mean u_{\parallel} of -0.59 GeV. This is important because a bias in u_{\parallel} would translate into a bias in the determination of m_T [Eq. (35)]. The agreement means that recoil momentum response and resolution and the u_{\parallel} efficiency parametrization describe the data well. Figures 48–50 show u_{\perp} , u_T , and the azimuthal difference between electron and recoil directions from Monte Carlo and W data. The Kolmogorov-Smirnov probabilities for Figs. 47–50 are $\kappa = 0.15, 0.38, 0.16,$ and 0.11 , respectively.

VIII. CONSTRAINTS ON THE W BOSON p_T SPECTRUM

A. Parameters

Since we cannot reconstruct a Lorentz-invariant mass for $W \rightarrow e\nu$ decays, knowledge of the transverse momentum dis-




FIG. 50. Azimuthal difference between electron and recoil directions for the W data (●) and the Monte Carlo simulation (solid line).




FIG. 51. Comparison of the $p_T(ee)$ data (●) and simulation (solid line) for the best fit g_2 using MRSA' parton distribution functions.

tribution of the W bosons is necessary to measure the mass from the kinematic distributions. Theoretical calculations provide a formalism to describe the boson p_T spectrum, but it includes the phenomenological parameters g_1 , g_2 , and g_3 , which need to be determined experimentally (Sec. V B). In addition, the boson p_T spectrum also depends on the choice of parton distribution functions and Λ_{QCD} .

We can measure the W boson p_T spectrum only indirectly by measuring \vec{u}_T , the p_T of all particles that recoil against the W boson. Momentum conservation requires the W boson p_T to be equal and opposite to \vec{u}_T . The precision of the \vec{u}_T measurement is insufficient, especially for small u_T , to constrain the W spectrum as tightly as is necessary for a precise W mass measurement.

We therefore have to find other data sets to constrain the model. The formalism that describes the p_T spectrum of the W bosons has to simultaneously describe the p_T spectrum of Z bosons and the dilepton p_T spectrum from Drell-Yan production with the same model parameter values. The authors of Ref. [30] find

$$\begin{aligned} g_1 &= 0.11^{+0.04}_{-0.03} \text{ GeV}^2, \\ g_2 &= 0.58^{+0.1}_{-0.2} \text{ GeV}^2, \\ g_3 &= -1.5^{+0.1}_{-0.1} \text{ GeV}^{-1} \end{aligned} \quad (50)$$

for the mass cutoff $Q_0 = 1.6$ GeV in Eq. (25) and CTEQ2M parton distribution functions, by fitting Drell-Yan and Z data at different values of Q^2 . We further constrain these parameters using our much larger Z data sample.

B. Determination of g_2 from $Z \rightarrow ee$ data

The p_T of Z bosons can be measured more precisely than the p_T of W bosons by using the e^+e^- pairs from their decays. Figure 51 shows the $p_T(ee)$ spectrum observed in the data.

To reduce the background contamination of the sample, the invariant mass of Z candidates must be within 10.5 GeV




FIG. 52. Background parametrizations for the $p_T(ee)$ spectrum.

of the Z peak position. This mass window requirement reduces the background fraction to 2.5%, as determined from the dielectron invariant mass spectrum. As such, it includes a contribution from Drell-Yan e^+e^- production, which has a p_T spectrum similar to the signal and should not be counted as background in this case. To account for this uncertainty we assign an error to the background fraction of $\pm 2.5\%$.

The shape of the background is fixed by a sample of events with two electromagnetic clusters which pass the same kinematic requirements as our $Z \rightarrow ee$ sample, but fail the electron identification cuts [47] (sample 1). As a cross-check, we also use events with two jets, each with more than 70% of its energy in the EM calorimeter (sample 2). Parametrizations of the two background shapes are shown in Fig. 52. Their difference is taken to be the uncertainty in the background shape.

We use the fast Monte Carlo model to predict the $p_T(ee)$ spectrum from $Z \rightarrow ee$ decays for different sets of parameter values. The fast Monte Carlo model simulates the detector acceptance and resolution as discussed in the previous sec-




FIG. 53. Predicted $p_T(ee)$ spectra after detector simulation using MRSA' parton distribution functions and $g_2 = 0.18, 0.58,$ and 0.98 GeV^2 .

TABLE V. Fitted values of g_2 for different parton distribution functions. Uncertainties are statistical only.

	$p_T(ee) < 15$ GeV (GeV ²)	$p_T(ee) < 30$ GeV (GeV ²)	$\Delta\phi(ee)$ (GeV ²)
MRSA'	0.59 ± 0.10	0.57 ± 0.10	0.64 ± 0.14
MRSD-'	0.61 ± 0.10	0.59 ± 0.10	0.70 ± 0.15
CTEQ3M	0.54 ± 0.10	0.52 ± 0.10	0.57 ± 0.13
CTEQ2M	0.61 ± 0.10	0.58 ± 0.10	0.67 ± 0.14

tions. Figure 53 shows the $p_T(ee)$ spectra predicted by the fast Monte Carlo model for the MRSA' parton distribution functions and three values of g_2 , with g_1 and g_3 fixed at the values given in Eq. (50).

The dominant effect of varying g_2 is to change the mean boson p_T . Properly normalized and with the background contribution added, we use these distributions as probability density functions to perform a maximum likelihood fit for g_2 . For a set of discrete values of g_2 , we compute the joint likelihood L of the observed $p_T(ee)$ spectrum. We then fit $\log L$ as a function of g_2 with a third-order polynomial. The maximum of the polynomial gives the fitted value of g_2 . The value of g_2 has to be fit independently for each parton distribution function choice. We perform fits for four choices of parton distribution functions: MRSA', MRSD-', CTEQ2M, and CTEQ3M. We fit the spectrum over the range $p_T(ee) < 15$ GeV, which corresponds to the range accepted by the W selection cuts. The fits describe the data well. Table V lists the fitted values for g_2 for the different parton distribution function choices. The result of the CTEQ2M fit is in good agreement with the value in Eq. (50).

We estimate systematic uncertainties in the g_2 fit by running the fast Monte Carlo model with different parameter values and refitting the predicted $p_T(ee)$ spectrum with the nominal probability density functions. The uncertainties in electron momentum response and resolution, $u_{||}$ efficiency parametrization, fiducial cuts, model of radiative decays, and background translate into a systematic uncertainty in g_2 of 0.05 GeV².

As cross-checks, we fit the $p_T(ee)$ spectrum for $p_T(ee) < 30$ GeV and the spectrum of the azimuthal separation $\Delta\phi(ee)$ between the two electrons to constrain g_2 . The $\Delta\phi(ee)$ spectrum has smaller systematic uncertainties, but less statistical sensitivity to g_2 than the $p_T(ee)$ spectrum. In Table V we also quote the results for g_2 from the fits to $p_T(ee) < 30$ GeV and the $\Delta\phi(ee)$ spectrum.

The Monte Carlo prediction for the fitted g_2 value using MRSA' parton distribution functions is superimposed as a smooth curve on Fig. 51. The Kolmogorov-Smirnov probability that the two distributions are from the same parent distribution is $\kappa=0.72$, and the χ^2 is 25.5 for 29 degrees of freedom. Both of these tests indicate a good fit. We use this model to compute the probability density functions for the final fits to the kinematic spectra from the W sample.

IX. BACKGROUNDS

A. $W \rightarrow \tau\nu \rightarrow e\nu\bar{\nu}\nu$

The decay $W \rightarrow \tau\nu \rightarrow e\nu\bar{\nu}\nu$ is topologically indistinguishable from $W \rightarrow e\nu$. It is included in the fast Monte Carlo




FIG. 54. p_T spectra of a sample of events passing electron identification cuts (solid line) and a sample of events failing the cuts (●).

simulation (Sec. V). This decay is suppressed by the branching fraction for $\tau \rightarrow e\nu\bar{\nu}$, $(17.83 \pm 0.08)\%$ [18], and by the lepton p_T cuts. It accounts for 1.6% of events in the W sample.

B. Hadronic background

QCD processes can fake the signature of a $W \rightarrow e\nu$ decay if a hadronic jet fakes the electron signature and the transverse momentum balance is mismeasured.

We estimate this background from the p_T spectrum of events with an electromagnetic cluster. Electromagnetic clusters in events with low p_T are almost all due to jets. A fraction satisfies our electron selection criteria and fakes an electron. From the shape of the p_T spectrum for these events, we determine how likely it is for these events to have sufficient p_T to enter our W sample.

We determine this shape by selecting isolated electromagnetic clusters that have $\chi^2 > 200$ and $\sigma_{\text{trk}} > 10$. Almost all electrons fail this cut, so that the remaining sample consists almost entirely of hadrons. We use data taken by a trigger without the p_T requirement to study the efficiencies of this cut for jets. For $p_T < 10$ GeV we find 1973 such events, while in the same sample 3674 satisfy our electron selection criteria. If we normalize the background spectrum to the electron sample, we obtain an estimate of the hadronic background in an electron candidate sample. Figure 54 shows the p_T spectra of both samples, normalized for $p_T < 10$ GeV.

In the data collected with the W trigger, we find 204 events that satisfy all the fiducial and kinematic cuts, listed in Sec. IV for the W sample, and have $\chi^2 > 200$ and $\sigma_{\text{trk}} > 10$. We therefore estimate that 374 background events entered the signal sample. This corresponds to a fraction of the total W sample after all cuts of $f_{\text{had}} = (1.3 \pm 0.2)\%$. For a looser cut on the recoil p_T , $u_T < 30$ GeV, we find $f_{\text{had}} = (1.6 \pm 0.3)\%$. The error is dominated by uncertainty in the relative normalization of the two samples at low p_T . Figure 55 shows the background fraction as a function of luminosity. There is no evidence for a significant luminosity dependence. We use the background events with $p_T(\nu) > 25$ GeV to esti-




FIG. 55. Fraction of hadron background as a function of luminosity.

mate the shape of the background contributions to the $p_T(e)$, $p_T(\nu)$, and m_T spectra (Fig. 56).

C. $Z \rightarrow ee$

To estimate the fraction of $Z \rightarrow ee$ events which satisfy the W selection, we use a Monte Carlo sample of approximately 100 000 $Z \rightarrow ee$ events generated with the HERWIG program and a detector simulation based on GEANT. The boson p_T spectrum generated by HERWIG agrees reasonably well with the calculation in Ref. [30]. $Z \rightarrow ee$ decays typically enter the W sample when one electron satisfies the W cuts and the second electron is lost or mismeasured, causing the event to have large \cancel{p}_T .

Approximately 1.1% of the $Z \rightarrow ee$ events have an electron with pseudorapidity $|\eta| > 4.0$, which is the acceptance limit of the end calorimeters. The fraction of $Z \rightarrow ee$ events which contain one electron with $|\eta(e_1)| < 1.0$ and $p_T(e) > 25$ GeV, and another with $|\eta(e_2)| > 4.0$, is approximately 0.04%. The contribution from the case of an electron lost through the beam pipe is therefore relatively small.

An electron is most frequently mismeasured when it goes into the regions between the CC and one of the ECs, which are covered only by the hadronic section of the calorimeter. These electrons therefore cannot be identified, and their energy is measured in the hadronic calorimeter. A large \cancel{p}_T is more likely for these events than when both electrons hit the EM calorimeters. The mismeasured electron contributes to the recoil when the event is treated as a W . The fraction of Z




FIG. 56. Shape of m_T , $p_T(e)$, and $p_T(\nu)$ spectra from hadron (solid line), Z (dashed line), and $\tau \rightarrow$ hadron backgrounds (dotted line) with the proper relative normalization.

events in the W sample therefore depends on the u_T cut.

We find that 10 987 Monte Carlo events pass the CC-CC $Z \rightarrow ee$ selection, and 758 (1318) pass the W selection with a recoil cut of 15 (30) GeV. The fraction of Z events in the W sample is therefore $f_Z = (0.42 \pm 0.08)\%$ for $u_T < 15$ GeV and $(0.62 \pm 0.08)\%$ for $u_T < 30$ GeV. The uncertainties quoted include systematic uncertainties in the matching of momentum scales between Monte Carlo and collider data. Figure 56 shows the distributions of $p_T(e)$, $p_T(\nu)$, and m_T for the events that satisfy the W selection.

D. $W \rightarrow \tau\nu \rightarrow$ hadrons+ X

We estimate the background due to $W \rightarrow \tau\nu$ followed by a hadronic τ decay based on two Monte Carlo samples. In a sample of $W \rightarrow \tau\nu \rightarrow$ hadrons+ X simulated using GEANT, 65 out of 4514 events pass the fiducial and kinematic cuts of the W sample. We use a sample of $W \rightarrow \tau\nu \rightarrow$ hadrons+ X simulated by replacing the electron shower in $W \rightarrow e\nu$ decays from collider data with the hadrons from a τ decay, generated by a Monte Carlo simulation, to estimate the probability of the τ decay products to fake an electron. Of 552 events that pass the fiducial and kinematic cuts, 145 pass the electron identification criteria. With the hadronic branching fraction for τ 's, $B(\tau \rightarrow$ hadrons) = 64%, we estimate a contamination of the W sample of 0.24% from hadronic τ decays. The expected background shapes are plotted in Fig. 56.

TABLE VI. Results of the Monte Carlo ensemble tests fitting the M_W mass for 105 samples of 28 323 events.

	Mean (GeV)	rms (GeV)	Correlation matrix		
			m_T	$p_T(e)$	$p_T(\nu)$
m_T	80.404	0.067	1	0.669	0.630
$p_T(e)$	80.415	0.091	0.669	1	0.180
$p_T(\nu)$	80.389	0.105	0.630	0.180	1




FIG. 57. Spectrum of $p_T(e)$ from the W data. The superimposed curve shows the maximum likelihood fit and the shaded region the estimated background.

E. Cosmic rays

Cosmic-ray muons can cause backgrounds when they coincide with a beam crossing and radiate a photon of sufficient energy to mimic the signature of the electron from $W \rightarrow e\nu$ decays. We measure this background by searching for muons near the electrons in the W signal sample. The muons have to be within 10° of the electron in azimuth. Using muon selection criteria similar to those in Ref. [49] we observe 18 events with such muons in the W sample. We estimate the fraction of cosmic-ray events in the W sample to be $(0.2 \pm 0.1)\%$. The effect of this background on the W mass measurement is negligible.

X. MASS FITS

A. Maximum likelihood fitting procedure

We use a binned maximum likelihood fit to extract the W mass. Using the fast Monte Carlo program, we compute the m_T , $p_T(e)$, and $p_T(\nu)$ spectra for 200 hypothesized values of the W mass between 79.4 and 81.4 GeV. For the m_T spectrum we use 100 MeV bins, and for the lepton p_T spectra we use 50 MeV bins. The statistical precision of the spectra for the W mass fit corresponds to about 4×10^6 W decays. When fitting the collider data spectra we add the background contributions with the shapes and normalizations described in Sec. IX to the signal spectra. We normalize the spectra within the fit interval and interpret them as probability density functions to compute the likelihood

$$L(m) = \prod_{i=1}^N p_i(m)^{n_i}, \quad (51)$$

where $p_i(m)$ is the probability density for bin i , assuming $M_W = m$, and n_i is the number of data entries in bin i . The product runs over all N bins inside the fit interval. We fit $-\ln[L(m)]$ with a quadratic function of m . The value of m at which the function assumes its minimum is the fitted value




FIG. 58. χ distribution for the fit to the $p_T(e)$ spectrum.

of the W mass, and the 68% confidence level interval is the interval in m for which $-\ln[L(m)]$ is within half a unit of its minimum.

As a consistency check of the fitting procedure, we generate 105 Monte Carlo ensembles of 28 323 events each with $M_W = 80.4$ GeV. We then fit these ensembles with the same probability density functions as the collider data spectra, except that we do not include the background contributions. Table VI lists the mean, rms, and correlation matrix of the fitted values.

B. Electron p_T spectrum

We fit the $p_T(e)$ spectrum in the region $30 < p_T(e) < 50$ GeV. There are 22 898 events in this interval. The data points in Fig. 57 represent the $p_T(e)$ spectrum from the W sample. The solid line shows the sum of the simulated W signal and the estimated background for the best fit, and the shaded region indicates the sum of the estimated hadronic, $Z \rightarrow ee$, and $W \rightarrow \tau\nu \rightarrow \text{hadrons} + X$ backgrounds. The maximum likelihood fit gives




FIG. 59. Variation of the fitted mass with the $p_T(e)$ fit window limits. See text for details.




FIG. 60. Spectrum of m_T from the W data. The superimposed curve shows the maximum likelihood fit, and the shaded region shows the estimated background.

$$M_W = 80.475 \pm 0.087 \text{ GeV} \quad (52)$$

for the W mass.

As a goodness-of-fit test, we divide the fit interval into 0.5 GeV bins, normalize the integral of the probability density function to the number of events in the fit interval, and compute $\chi^2 = \sum_{i=1}^N (y_i - P_i)^2 / y_i$. The sum runs over all N bins, y_i is the observed number of events in bin i , and P_i is the integral of the normalized probability density function over bin i . The parent distribution is the χ^2 distribution for $N - 2$ degrees of freedom. For the spectra in Fig. 57 we compute $\chi^2 = 40.6$. For 40 bins there is a 35% probability for $\chi^2 \geq 40.6$. Figure 58 shows the contributions $\chi_i = (y_i - P_i) / \sqrt{y_i}$ to χ^2 for the 40 bins in the fit interval.

We also compare the observed spectrum to the probability density function using the Kolmogorov-Smirnov test. For a comparison within the fit window, we obtain $\kappa = 0.81$ and for the entire histogram $\kappa = 0.83$.

Figure 59 shows the sensitivity of the fitted mass value to the choice of fit interval. The points in the two plots indicate the observed deviation of the fitted mass from the value




FIG. 61. Likelihood function for the m_T fit.




FIG. 62. χ distribution for the fit to the m_T spectrum.

given in Eq. (52). We expect some variation due to statistical fluctuations in the spectrum and systematic uncertainties in the probability density functions. We estimate the effect due to statistical fluctuations using the Monte Carlo ensembles described above. We expect the fitted values to be inside the shaded regions indicated in the two plots with 68% probability. The dashed lines indicate the statistical error for the nominal fit.

All tests show that the probability density function provides a good description of the observed spectrum.

C. Transverse mass spectrum

Figure 60 shows the m_T spectrum. The points are the observed spectrum, the solid line shows signal plus background for the best fit, and the shaded region indicates the estimated background contamination. We fit in the interval $60 < m_T < 90$ GeV. There are 23 068 events in this interval. Figure 61 shows $-\ln[L(m)/L_0]$ for this fit where L_0 is an arbitrary number. The best fit occurs for

$$M_W = 80.438 \pm 0.070 \text{ GeV}. \quad (53)$$




FIG. 63. Variation of the fitted mass with the m_T fit window limits. See text for details.




FIG. 64. Spectrum of $p_T(\nu)$ from the W data. The superimposed curve shows the maximum likelihood fit, and the shaded region shows the estimated background.

Figure 62 shows the deviation of the data from the fit. Summing over all bins in the fitting window, we get $\chi^2 = 79.5$ for 60 bins. For 60 bins there is a 3% probability to obtain a larger value. The Kolmogorov-Smirnov test gives $\kappa = 0.25$ within the fit window and $\kappa = 0.84$ for the entire histogram. Figure 63 shows the sensitivity of the fitted mass to the choice of fit interval.

In spite of the somewhat large value of χ^2 , there is no structure apparent in Fig. 62 that would indicate that there is a systematic difference between the shapes of the observed spectrum and the probability density function. The large χ^2 can be attributed to a few bins that are scattered over the entire fit interval, indicating statistical fluctuations in the data. This is consistent with the good Kolmogorov-Smirnov probability which is more sensitive to the shape of the distribution and insensitive to the binning.

XI. CONSISTENCY CHECKS

A. Neutrino p_T spectrum

As a consistency check, we also fit the $p_T(\nu)$ spectrum, although this measurement is subject to much larger system-




FIG. 65. χ distribution for the fit to the $p_T(\nu)$ spectrum.




FIG. 66. Fitted W boson masses (a) in bins of luminosity from the m_T (\bullet), $p_T(e)$ (\circ), and $p_T(\nu)$ ($*$) fits (the points are offset for clarity) and the fitted Z boson masses (b). The solid line is the central value for the m_T and $m(ee)$ mass fits, respectively, over the entire luminosity range, and the dashed lines are the statistical errors.

atic uncertainties than the m_T and $p_T(e)$ fits. Figure 64 shows the observed spectrum (points), signal plus background for the best fit (solid line), and the estimated background (shaded region). For the fit interval $30 < p_T(\nu) < 50$ GeV, the fitted mass is $M_W = 80.37 \pm 0.11$ GeV, in good agreement with the m_T and $p_T(e)$ fits. We compute $\chi^2 = 31.8$. The probability for a larger value is 75%. The Kolmogorov-Smirnov test gives $\kappa = 0.20$ within the fit window and $\kappa = 0.69$ for the entire histogram. Figure 65 shows the deviation χ between data and fit. There is an indication of a systematic deviation between the observed spectrum and the resolution function. This effect is not very significant. For example, when we increase the hadronic resolution parameter α_{mb} in the simulation to 1.11, which corresponds to about 1.5 standard deviations, this indication of a deviation between data and Monte Carlo simulation vanishes.

B. Luminosity dependence

We divide the W and Z data samples into four luminosity bins

$$\mathcal{L} \leq 5 \times 10^{30} \text{ cm}^{-2} \text{ s}^{-1},$$

$$5 \times 10^{30} < \mathcal{L} \leq 7 \times 10^{30} \text{ cm}^{-2} \text{ s}^{-1},$$

$$7 \times 10^{30} < \mathcal{L} \leq 9 \times 10^{30} \text{ cm}^{-2} \text{ s}^{-1},$$

$$\mathcal{L} > 9 \times 10^{30} \text{ cm}^{-2} \text{ s}^{-1},$$

and generate resolution functions for the luminosity distribu-




FIG. 67. Comparison of the m_T spectra between the data (\bullet) and Monte Carlo simulation (solid line) for the nominal mass of 80.44 GeV in bins of u_T . Starting from the top, the data are selected with $u_T < 5$ GeV, $5 < u_T < 10$ GeV, and $10 < u_T < 15$ GeV.

tion of these four subsamples. We fit the transverse mass and lepton p_T spectra from the W samples and the dielectron invariant mass spectra from the Z samples in each bin. The fitted masses are plotted in Fig. 66. The errors are statistical only. We compute the χ^2 with respect to the W mass fit to the m_T spectrum from the entire data sample. The χ^2 per degree of freedom (dof) for the $p_T(e)$ fit is 1.9/4 and for the $p_T(\nu)$ fit is 2.4/4. The m_T fit has a χ^2/dof of 2.7/3. The solid and dashed lines in the top plot indicate the W mass value and statistical uncertainty from the fit to the m_T spectrum of the entire data sample. All measurements are in very good agreement with this value. In the bottom plot the lines indicate the Z mass fit to the $m(ee)$ spectrum of the entire Z data sample. The measurements in the four luminosity bins have a χ^2/dof of 1.0/3.

C. Dependence on the u_T cut

We change the cuts on the recoil momentum u_T and study how well the fast Monte Carlo simulation reproduces the




FIG. 68. Spectra of m_T from W data with $u_{\parallel} < 0$ (\circ) and $u_{\parallel} > 0$ (\bullet) compared to Monte Carlo simulations (solid line).




FIG. 69. Spectra of $p_T(e)$ from W data with $u_{\parallel} < 0$ (\circ) and $u_{\parallel} > 0$ (\bullet) compared to Monte Carlo simulations (solid line).

variations in the spectra. We split the W sample into subsamples with $u_{\parallel} > 0$ GeV, $u_{\parallel} < 0$ GeV, $u_T < 5$ GeV, $5 < u_T < 10$ GeV, and $10 < u_T < 15$ GeV. In the simulation we fix the W mass to the value from the m_T fit in Eq. (53). Figure 67 shows the m_T spectra from the collider data and Monte Carlo simulation for the $u_T < 5$ GeV, $5 < u_T < 10$ GeV, and $10 < u_T < 15$ GeV subsamples. Figures 68–70 show the m_T , $p_T(e)$, and $p_T(\nu)$ spectra from the collider data for the subsamples with $u_{\parallel} > 0$ and $u_{\parallel} < 0$ and the corresponding Monte Carlo predictions. Table VII lists the results of comparisons of collider data and Monte Carlo spectra using the Kolmogorov-Smirnov test. Although there is significant variation among the shapes of the spectra for the different cuts, the fast Monte Carlo simulation models them well. Table VII also lists the results of comparisons of collider data and Monte Carlo spectra for a W sample selected with $u_T < 30$ GeV which consists of 32 361 events.

D. Dependence on fiducial cuts

We divide the azimuth of the recoil momentum, $\phi(R)$, into eight bins. This binning is sensitive to azimuthal non-




FIG. 70. Spectra of $p_T(\nu)$ from W data with $u_{\parallel} < 0$ (\circ) and $u_{\parallel} > 0$ (\bullet) compared to Monte Carlo simulations (solid line).

TABLE VII. Confidence levels from Kolmogorov-Smirnov tests comparing collider data to Monte Carlo predictions for $M_W = 80.44$ GeV.

Interval	m_T	$p_T(e)$	$p_T(\nu)$
	60–90 GeV	30–50 GeV	30–50 GeV
$u_T < 15$ GeV	0.25	0.81	0.20
$u_T < 5$ GeV	0.01	0.03	0.03
$5 \leq u_T < 10$ GeV	0.17	0.83	0.21
$10 \leq u_T < 15$ GeV	0.85	0.68	0.69
$u_T < 30$ GeV	0.55	0.99	0.58
$u_{\parallel} < 0$	0.19	0.78	0.25
$u_{\parallel} > 0$	0.61	0.80	0.48
Interval	50–100 GeV	25–55 GeV	25–55 GeV
	$u_T < 15$ GeV	0.84	0.83
$u_T < 5$ GeV	10^{-3}	2×10^{-3}	0.01
$5 \leq u_T < 10$ GeV	0.10	0.44	0.29
$10 \leq u_T < 15$ GeV	0.61	0.98	0.31
$u_T < 30$ GeV	0.92	0.80	0.28
$u_{\parallel} < 0$	0.77	0.67	0.62
$u_{\parallel} > 0$	0.60	0.66	0.73

uniformities in the recoil momentum measurement, e.g., because of background from the Main Ring. Figure 71 shows the fitted W mass values versus $\phi(R)$. The Main Ring is located at $\phi \sim \pi/2$, and any biases caused by background from the Main Ring should appear as structure in this direction or in the opposite direction. The rms of the eight data points is 124 MeV, consistent with the statistical uncertainty of 200 MeV for the data points. Thus the data are consistent with azimuthal uniformity.

We divide the azimuthal direction of the electron, $\phi(e)$, into 32 bins corresponding to the 32 azimuthal modules of the CC-EM section. Figure 72 shows the fitted W mass values versus $\phi(e)$. The statistical uncertainty of the data points is 400 MeV, and the rms of the 32 points is 600 MeV. Thus there is a 0.6% nonuniformity in the response of the CC-EM




FIG. 71. Variation in the W mass from the m_T fit as a function of $\phi(R)$.




FIG. 72. Variation in the W mass from the m_T fit as a function of $\phi(e)$.

section, consistent with the module-to-module calibration of 0.5% [25].

Finally, we fit the m_T spectrum from the W sample and the $m(ee)$ spectrum from the Z sample for different pseudorapidity cuts on the electron direction. We use cuts of $|\eta(e)| < 1.0, 0.7, 0.5,$ and 0.3 . Figure 73 shows the change in the W mass versus the $\eta(e)$ cut using the electron energy scale calibration from the corresponding Z sample. The shaded region indicates the statistical error. Within the uncertainties the mass is independent of the $\eta(e)$ cut.

XII. UNCERTAINTIES IN THE MEASUREMENT

A. Statistical uncertainties

Table VIII lists the uncertainties, rounded to the nearest 5 MeV, in the W measurement due to the finite sizes of the W and Z samples used in the fits to the m_T , $p_T(e)$, $p_T(\nu)$, and $m(ee)$ spectra. The statistical uncertainty due to the finite Z sample propagates into the W mass measurement through the electron energy scale α_{EM} .




FIG. 73. Variation in the W mass versus the $\eta(e)$ cut. The shaded region is the expected statistical variation.

TABLE VIII. Uncertainties in the W mass measurement due to finite sample sizes.

	m_T fit (MeV)	$p_T(e)$ fit (MeV)	$p_T(\nu)$ fit (MeV)
W sample	70	85	105
Z sample	65	65	65
Total	95	105	125

B. W production and decay model

1. Sources of uncertainty

Uncertainties in the W production and decay model arise from the following sources: the phenomenological parameters in the calculation of the $p_T(W)$ spectrum, the choice of parton distribution functions, radiative decays, and the W boson width. In the following we describe how we assess the size of the systematic uncertainties introduced by each of these. We summarize the size of the uncertainties in Table IX, rounded to the nearest 5 MeV.

2. W boson p_T spectrum


In Sec. VIII we determine g_2 so that the predicted $p_T(ee)$ spectrum agrees with the Z data. In order to quantify the uncertainty in the boson p_T spectra, we need to consider variations in all four parameters Λ_{QCD} , g_1 , g_2 , and g_3 . We use a series of modified CTEQ3M parton distribution functions fit with Λ_{QCD} fixed at discrete values [48] to study the variations in the $p_T(ee)$ spectrum and the fitted W boson mass with these parameters.

We cannot constrain all these four parameters simultaneously by using only our Z data. We therefore introduce an external constraint on Λ_{QCD} . The CTEQ3M fits prefer $\Lambda_{\text{QCD}}=158$ MeV, but are also consistent with somewhat higher values [42]. Other measurements give a combined value of $\Lambda_{\text{QCD}}=209^{+39}_{-33}$ MeV [18]. All data are consistent with Λ_{QCD} between 150 and 250 MeV, which we use as the range over which Λ_{QCD} is allowed to vary.

The requirement that the fast Monte Carlo prediction for the average $p_T(ee)$ over the range $p_T(ee)<15$ GeV, corrected for background contributions, must agree with the value observed in the Z data, $\langle p_T(ee) \rangle = 6.05 \pm 0.07$ GeV, couples the values of Λ_{QCD} and g_2 . Figure 74 shows a plot of g_2 versus Λ_{QCD} . For any pair of values on the curve, the

TABLE IX. Uncertainties in the W mass measurement due to W production and decay model.

	m_T fit (MeV)	$p_T(e)$ fit (MeV)	$p_T(\nu)$ fit (MeV)
$p_T(W)$ spectrum	10	50	25
Parton distribution functions	20	50	30
Parton luminosity β	10	10	10
Radiative decays	15	15	15
W width	10	10	10
Total	30	75	45

FIG. 74. Value of g_2 as a function of Λ_{QCD} . The error bar indicates the uncertainty in g_2 for fixed Λ_{QCD} .

fast Monte Carlo model predicts a value of $\langle p_T(ee) \rangle$ that agrees with the Z data. For any fixed value of Λ_{QCD} , g_2 is determined to a precision of 0.12 GeV^2 . This error includes the statistical uncertainty (0.09 GeV^2) and the systematic uncertainty due to normalization and shape of the background (0.07 GeV^2). All other uncertainties, e.g., due to electron momentum resolution and response or selection biases, are negligible.

If we fix Λ_{QCD} and g_2 , the requirement that the average $p_T(ee)$ predicted by the fast Monte Carlo model agree with the data allows an additional variation in the parameters g_1 and g_3 . The residual uncertainty in the measured W boson mass due to this variation, however, is small compared to the uncertainty due to the variation allowed in g_2 and Λ_{QCD} , and we neglect it. Finally, we obtain the uncertainties in the fitted W boson mass listed in Table IX.

3. Parton distribution functions

The choice of parton distribution function used to describe the momentum distribution of the constituents of the proton and antiproton affects several components of the model: the parton luminosity slope β , and the rapidity and transverse momentum spectrum of the W .

Using several modern parton distribution function sets as input to the fast Monte Carlo model, we generate m_T and lepton p_T spectra. In each case we use the value of g_2 measured for that parton distribution function set using our Z data (Sec. VIII). We then fit them in the same way as the

TABLE X. Variation of fitted W mass with choice of parton distribution function.

	m_T fit (MeV)	$p_T(e)$ fit (MeV)	$p_T(\nu)$ fit (MeV)
MRSA'	0	0	0
MRSD-'	20	19	20
CTEQ3M	5	48	22
CTEQ2M	-21	-17	-30

TABLE XI. Changes in fitted W and Z masses if radiative effects are varied.

Variation	m_T fit (MeV)	$p_T(e)$ fit (MeV)	$p_T(\nu)$ fit (MeV)	$m(ee)$ fit (MeV)
No radiative effects	50	43	30	143
Vary R_0 by ± 0.1	3	4	0	19

spectra from collider data, i.e., using MRSA' parton distribution functions. Table X lists the variation of the fitted W mass values relative to MRSA'.

The MRSA' and CTEQ3M parton distribution functions use the measured W charge asymmetry in $p\bar{p}$ collisions [50] as input to the fit. MRSD-' and CTEQ2M do not explicitly use the asymmetry. The asymmetry predicted by MRSD-' agrees with the measurement; that of CTEQ2M disagrees at the level of four standard deviations. We include CTEQ2M in our estimate of the uncertainty to provide an estimate of the possible variations with a rather large deviation from the measured asymmetry.

4. Parton luminosity

The uncertainty of 10^{-3} GeV $^{-1}$ in the parton luminosity slope β (Sec. V) translates into an uncertainty in the fitted W mass. We estimate the sensitivity in the fitted W mass by fitting Monte Carlo spectra generated with different values of β .

5. Radiative decays

We assign an error to the modeling of radiative decays based on varying the detector parameters E_0 and R_0 (Sec. V). E_0 defines the minimum photon energy generated and corresponds to a cutoff below which the photon does not reach the calorimeter. R_0 defines the maximum separation between the photon and electron directions above which the photon energy is not included in the electron shower. In general, radiation shifts the fitted mass down for the transverse mass and electron fits, because for a fraction of the events the photon energy is subtracted from the electron. Hence increasing R_0 decreases the radiative shift. Similarly, decreasing E_0 decreases the radiative shift. Both the fitted W and Z masses depend on these parameters. Table XI lists the change in the fitted masses if radiative effects are turned off completely. To estimate the systematic error, we fit Monte Carlo spectra generated with different values for E_0 and R_0 . For the low value of $E_0=50$ MeV that we use in the simulation, the dependence of the fits on this parameter is negligible. The changes in the mass fits when varying R_0 by ± 0.1 are also listed in Table XI. After propagating the change in the Z mass into the electron response, the result of the W mass measurement changes by about 15 MeV for all three spectra.

There are also theoretical uncertainties in the radiative decay calculation. Initial-state QED radiation is not included in the calculation of Ref. [41]. However, initial-state radiation does not affect the kinematic distributions used to fit the mass in the final state. Finally, the calculation includes only

TABLE XII. Uncertainties in the W mass measurement due to detector model parameters.

	m_T fit (MeV)	$p_T(e)$ fit (MeV)	$p_T(\nu)$ fit (MeV)
Calorimeter linearity	20	20	20
Calorimeter uniformity	10	10	10
Electron resolution	25	15	30
Electron angle calibration	30	30	30
Electron removal	15	15	20
Selection bias	5	10	20
Recoil resolution	25	10	90
Recoil response	20	15	45
Total	60	50	115

processes in which a single photon is radiated. We use the code provided by the authors of Ref. [51] to estimate the shift introduced in the measured W mass by neglecting two-photon emission. We find that two photons, with $p_T > 100$ MeV and separated by $\Delta R > 0.3$ from the electron, are radiated in about 0.24% of all $W \rightarrow e\nu$ decays. This reduces the mean value of m_T within the fit window by 3 MeV. In 1.1% of all $Z \rightarrow ee$ decays, two photons, with $p_T > 100$ MeV and separated by $\Delta R > 0.3$ from the electrons, are radiated. We add the dielectron mass spectrum of these $Z \rightarrow ee\gamma\gamma$ events to our simulated Z boson line shape and fit the modified line shape. The fitted mass decreases by 10 MeV. This shift requires an adjustment of the energy scale calibration factor α_{EM} by 10^{-4} . Neglecting two-photon emission in both W and Z boson decays then increases the measured W mass by about 5 MeV. Since this effect is an order of magnitude smaller than the statistical uncertainty in our measurement, we do not correct for it, but add it in quadrature to the uncertainty due to radiative corrections.

6. W boson width

To determine the sensitivity of the fitted W mass to the W width, we generate m_T and lepton p_T spectra using the fast Monte Carlo model with a range of widths and fit them with the nominal templates. The uncertainty in the fitted W mass corresponds to the uncertainty in the measured value of $\Gamma_W = 2.062 \pm 0.059$ GeV [36].

C. Detector model parameters

The uncertainties in the parameters of the detector model determined in Secs. VI–VII translate into uncertainties in the W mass measurement. We study the sensitivity of the W mass measurement to the values of the parameters by fitting the data with spectra generated by the fast Monte Carlo model with modified input parameters.

Table XII lists the uncertainties in the measured W mass, caused by the individual parameters. We assign sets of correlated parameters to the same item in the table. Correlations between items are negligible. For each item the uncertainty is determined to typically 5 MeV for the m_T fit and 10 MeV for the lepton p_T fits. We therefore round them to the nearest

TABLE XIII. Uncertainties in the W mass measurement due to backgrounds.

	m_T fit (MeV)	$p_T(e)$ fit (MeV)	$p_T(\nu)$ fit (MeV)
Hadrons	10	15	20
$Z \rightarrow ee$	5	10	5
$W \rightarrow \tau\nu$		negligible	
Cosmic rays		negligible	
Total	10	20	20

5 MeV in the table. To achieve this precision (10–20) $\times 10^6$ $W \rightarrow e\nu$ decays are simulated for each item.

The residual calorimeter nonlinearity is parametrized by the offset δ_{EM} . Calorimeter uniformity refers to a possible nonuniformity in response as a function of η . It is limited by the test beam data [12]. The electron momentum resolution is parametrized by c_{EM} . The electron angle calibration includes the effects of the parameters α_{CDC} and α_{CC} , discussed in Appendixes A and B. The recoil resolution is parametrized by α_{mb} and s_{rec} and the response by α_{rec} and β_{rec} . Electron removal refers to the bias Δu_{\parallel} introduced in the u_{\parallel} measurement by the removal of the cells occupied by the electron. Selection bias refers to the u_{\parallel} efficiency.

D. Backgrounds

We determine the sensitivity of the fit results to the assumed background normalizations and shapes by repeating the fits to the data with varied background shapes and normalizations. Table XIII lists the uncertainties introduced in the W boson mass measurement rounded to the nearest 5 MeV. If we fit the spectra without including the backgrounds in the probability density functions, the measurements would

change by -75 MeV (m_T spectrum), -80 MeV ($p_T(e)$ spectrum), and -130 MeV ($p_T(\nu)$ spectrum). These shifts are dominated by the $W \rightarrow \tau\nu \rightarrow e\nu\bar{\nu}$ background.

XIII. RESULTS

We present a precision measurement of the mass of the W boson. From a fit to the transverse mass spectrum, we measure

$$M_W = 80.44 \pm 0.10(\text{stat}) \pm 0.07(\text{syst}) \text{ GeV}. \quad (54)$$

Adding all errors in quadrature gives 115 MeV. Since we calibrate the electron energy scale against the known Z mass, we effectively measure the W and Z mass ratio

$$\frac{M_W}{M_Z} = 0.8821 \pm 0.0011(\text{stat}) \pm 0.0008(\text{syst}). \quad (55)$$

A fit to the transverse momentum spectrum of the decay electrons gives

$$M_W = 80.48 \pm 0.11(\text{stat}) \pm 0.09(\text{syst}) \text{ GeV}. \quad (56)$$

Adding all errors in quadrature gives 140 MeV. As expected, the measurement from the m_T spectrum has a larger uncertainty from detector effects (65 MeV) than that from the $p_T(e)$ spectrum (50 MeV). On the other hand, the m_T fit is less sensitive to the W production model (30 MeV) than the $p_T(e)$ fit (75 MeV). The good agreement between the two results indicates that we understand the ingredients of our model and their uncertainties. In the end the m_T fit gives the more precise result and we quote this as our final result. However, the fit to the $p_T(e)$ spectrum may become more competitive in the future with larger data samples and better constraints on the W production dynamics.

TABLE XIV. Summary of results from the 1992–1993 and 1994–1995 data sets with the common and uncorrelated errors.

	1992–1993	1994–1995	Common
M_W from m_T fit	80.35 GeV	80.44 GeV	
W statistics	140 MeV	70 MeV	
Z statistics	160 MeV	65 MeV	
Calorimeter linearity			20 MeV
Calorimeter uniformity			10 MeV
Electron resolution	70 MeV	20 MeV	
Electron angle calibration			30 MeV
Recoil resolution	90 MeV	25 MeV	
Recoil response	50 MeV	20 MeV	
Electron removal	35 MeV	15 MeV	
Selection bias	30 MeV	5 MeV	
Backgrounds	35 MeV	10 MeV	
W production/decay			30 MeV
Total uncertainty	255 MeV	105 MeV	50 MeV

TABLE XV. Previously published measurements of the W boson mass.

Measurement	M_W (GeV)	Reference
CDF 90	79.91 ± 0.39	[52]
UA2 92	80.36 ± 0.37	[10]
CDF 95	80.41 ± 0.18	[11]
D0 96	80.35 ± 0.27	[12]
L3 97	80.75 ± 0.27	[13]
ALEPH 97	80.80 ± 0.34	[14]
OPAL 97	80.31 ± 0.25	[15]
DELPHI 97	80.33 ± 0.31	[16]

Table XIV lists the D0 W mass measurements from fits to the m_T spectra from the 1992–1993 [12] and the 1994–1995 data sets and their uncertainties. As indicated in Table XIV, some errors are common to the two measurements. Since both analyses use the same W production and decay model, we assign the uncertainties quoted in Sec. XII B to both measurements. The precision of the electron angle calibration has improved compared to Ref. [12], and we use the reduced uncertainty for both measurements. All uncertainties due to detector model parameters, which were measured using statistically independent data sets, are uncorrelated because their precision is dominated by statistical fluctuations. In order to combine the two measurements, we weight them by their uncorrelated errors δ_a and δ_b :

$$M_W = \frac{M_a / \delta_a^2 + M_b / \delta_b^2}{1/\delta_a^2 + 1/\delta_b^2}. \quad (57)$$

The uncertainty is then given by




FIG. 75. Comparison of this measurement with previously published W mass measurements (Table XV). The shaded region indicates the predicted W mass value from global fits to the Z line shape data [9].




FIG. 76. Comparison of the W and top quark mass measurements by the D0 Collaboration with the standard model predictions for different Higgs boson masses [53]. The width of the bands for each Higgs boson mass value indicates the uncertainty due to the error in $\alpha(M_Z^2)$. Also shown is the range allowed by the MSSM [21].

$$\delta M_W = \sqrt{\frac{1}{1/\delta_a^2 + 1/\delta_b^2} + \delta^2}, \quad (58)$$

where δ is the common uncertainty from the third column in Table XIV. The combination of the D0 measurements from the 1992–1993 and 1994–1995 data gives

$$M_W = 80.43 \pm 0.11 \text{ GeV}. \quad (59)$$

The D0 measurement is in good agreement with previous measurements and is more precise than all the previously published measurements combined. Table XV lists previously published measurements with uncertainties below 500 MeV. A global fit to all electroweak measurements from the LEP experiments predicts $M_W = 80.329 \pm 0.041$ GeV [9]. Figure 75 gives a graphical representation of these data.

We evaluate the radiative corrections Δr , defined in Eq. (1). Our measurement of M_W from Eq. (59) leads to

$$\Delta r = -0.0288 \pm 0.0070, \quad (60)$$

4.1 standard deviations from the tree-level value. In Fig. 76




FIG. 77. Difference between the predicted and actual z positions of the track center of gravity.




FIG. 78. χ^2 versus α_{CDC} value. The arrows indicate the statistical error in the fit.

we compare the measured W and top quark masses [20] to the values predicted by the standard model for a range of Higgs boson mass values [53]. Also shown is the prediction from the calculation in Ref. [21] for a model involving supersymmetric particles assuming the chargino, Higgs boson, and left-handed selectron masses are greater than 90 GeV. The measured values are in agreement with the prediction of the standard model.

ACKNOWLEDGMENTS

We wish to thank U. Baur for helpful discussions. We thank the staffs at Fermilab and collaborating institutions for their contributions to this work, and acknowledge support from the Department of Energy and National Science Foundation (U.S.), Commissariat à l’Energie Atomique (France), State Committee for Science and Technology and Ministry for Atomic Energy (Russia), CNPq (Brazil), Departments of Atomic Energy and Science and Education (India), Colciencias (Colombia), CONACyT (Mexico), Ministry of Education and KOSEF (Korea), CONICET and UBACyT (Argentina), and CAPES (Brazil).

APPENDIX A: TRACK POSITION CALIBRATION

We use cosmic-ray muons which traverse the entire detector and pass close to the beam position to calibrate the z measurement of the track in the CDC. We predict the trajectory of the muon through the central detector by connecting the incoming and outgoing hits in the innermost muon chambers by a straight line. The centers of gravity of the incoming and outgoing CDC tracks are then calibrated relative to this line. Figure 77 shows the difference between the predicted and the actual z positions of the track centers of gravity. These data are fit to a straight line. We find the track position must be scaled by the fitted slope, $\alpha_{\text{CDC}} = 0.9868 \pm 0.0004$.

We also use a sample of low- p_T dimuon events from $p\bar{p}$ collisions where both muons originate from the same interaction vertex. We reconstruct the muon trajectories from their hits in the innermost muon chambers and the CDC. For

both muons we determine the point of closest approach of the trajectory to the beam, $z_{\text{vtx}}(\mu)$. We then scale the z position of the CDC track to minimize

$$\chi^2 = \sum_{\text{events}} \left(\frac{z_{\text{vtx}}(\mu_1) - z_{\text{vtx}}(\mu_2)}{\sigma_\mu} \right)^2, \quad (\text{A1})$$

where σ_μ is chosen so that the minimum value of χ^2 equals the number of events minus 1. The minimum occurs at $\alpha_{\text{CDC}} = 0.9863 \pm 0.0011$. The same analysis applied to a $Z \rightarrow \mu\mu$ sample gives $\alpha_{\text{CDC}} = 0.9878 \pm 0.0014$ and is shown in Fig. 78.

A scintillating fiber detector was inserted between the CDC and CC to calibrate the track z position. The detector is built from 20 modules, each constructed on an aluminum support plate 93.4 cm long and 16.5 cm wide. Scintillating fibers, 12.7 cm long, were laid across the width of the module every 11.43 cm along the support plate. The eight scintillating fibers on each module were connected to a clear waveguide and read out with a photomultiplier tube. The modules are mounted lengthwise along the cylinder of the CDC with half of the modules covering $+z$ and the other half $-z$. In the r - ϕ view each module subtends $\pi/16$ radians with the fibers running azimuthally. Because of spatial constraints, the entire CDC was not covered.

When a fiber is hit by a charged particle, the z position of the associated track, at the fiber radius, is compared with the fiber z position. The z position of the track at the radial position of the fiber is determined from the direction and center of gravity of the track. By comparing the z position of the track and the hit fiber, we determine that a scale of $\alpha_{\text{CDC}} = 0.989 \pm 0.001$ is needed to correct the track.

Combining all measurements of α_{CDC} gives $\alpha_{\text{CDC}} = 0.988 \pm 0.001$, which we use in the reconstruction of the electrons in the W and Z data samples.

APPENDIX B: ELECTRON SHOWER POSITION ALGORITHM

We determine the position of the electron shower centroid $\vec{x}_{\text{cal}} = (x_{\text{cal}}, y_{\text{cal}}, z_{\text{cal}})$ in the calorimeter from the energy depositions in the third EM layer by computing the weighted mean of the positions \vec{x}_i of the cell centers:

$$\vec{x}_{\text{cal}} = \frac{\sum_i w_i \vec{x}_i}{\sum_i w_i}. \quad (\text{B1})$$

The weights are given by

$$w_i = \max\left(0, w_0 + \log\left(\frac{E_i}{E(e)}\right)\right), \quad (\text{B2})$$

where E_i is the energy in cell i , w_0 is a parameter which depends upon $\eta(e)$, and $E(e)$ is the energy of the electron. We calibrate the algorithm using Monte Carlo electrons simulated using GEANT and electrons from the $Z \rightarrow ee$ data. We apply a polynomial correction as a function of z_{cal} and $\theta(e)$ based on the Monte Carlo electrons. We refine the calibration with the $Z \rightarrow ee$ data by exploiting the fact that both

electrons originate from the same vertex. Using the algorithm given by Eq. (A1), we determine a vertex for each electron from the shower centroid and the track center of gravity. We minimize the difference between the two vertex

positions as a function of a scale factor α_{CC} . More complex correction functions do not improve the χ^2 . The correction factor is $\alpha_{CC}=0.9980\pm 0.0005$, where the error includes possible variations of the functional form of the correction.

-
- [1] I. Adam, Ph.D. thesis, Columbia University, 1997, Nevis Report No. 294, http://www-d0.fnal.gov/publications_talks/thesis/adam/ian_thesis_all.ps.
- [2] E. Flattum, Ph.D. thesis, Michigan State University, 1996, http://www-d0.fnal.gov/publications_talks/thesis/flattum/eric_thesis.ps.
- [3] D0 Collaboration, B. Abbott *et al.*, Phys. Rev. Lett. **80**, 3008 (1998).
- [4] UA1 Collaboration, G. Arnison *et al.*, Phys. Lett. **122B**, 103 (1983).
- [5] UA2 Collaboration, M. Banner *et al.*, Phys. Lett. **122B**, 476 (1983).
- [6] UA1 Collaboration, G. Arnison *et al.*, Phys. Lett. **126B**, 398 (1983).
- [7] UA2 Collaboration, P. Bagnaia *et al.*, Phys. Lett. **129B**, 130 (1983).
- [8] S. L. Glashow, Nucl. Phys. **22**, 579 (1961); S. Weinberg, Phys. Rev. Lett. **19**, 1264 (1967); A. Salam, in *Relativistic Groups and Analyticity (Nobel Symposium No. 8)*, edited by N. Svartholm (Almqvist and Wiksell, Stockholm, 1968), p. 367.
- [9] The LEP Collaborations, the LEP Electroweak Working Group, and the SLD Heavy Flavour Group, Report No. CERN-PPE/97-154 (unpublished).
- [10] UA2 Collaboration, J. Alitti *et al.*, Phys. Lett. B **276**, 354 (1992). The value quoted in Table XV uses M_Z from Eq. (3).
- [11] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **75**, 11 (1995); Phys. Rev. D **52**, 4784 (1995).
- [12] D0 Collaboration, S. Abachi *et al.*, Phys. Rev. Lett. **77**, 3309 (1996); D0 Collaboration, B. Abbott *et al.* Phys. Rev. D **58**, 012 002 (1998).
- [13] L3 Collaboration, M. Acciarri *et al.*, Phys. Lett. B **413**, 176 (1997).
- [14] ALEPH Collaboration, R. Barate *et al.*, Phys. Lett. B **422**, 384 (1998).
- [15] OPAL Collaboration, K. Ackerstaff *et al.*, Eur. Phys. J. C **1**, 395–424 (1998).
- [16] Delphi Collaboration, P. Abreu *et al.*, Eur. Phys. J. C. **2**, 581 (1998).
- [17] A. Sirlin, Phys. Rev. D **22**, 971 (1980); W. Marciano and A. Sirlin, *ibid.* **22**, 2695 (1980); **31**, 213E (1985).
- [18] R. M. Barnett *et al.*, Phys. Rev. D **54**, 1 (1996).
- [19] S. Eidelmann and F. Jegerlehner, Z. Phys. C **67**, 585 (1995).
- [20] D0 Collaboration, S. Abachi *et al.*, Phys. Rev. Lett. **79**, 1197 (1997) and references therein.
- [21] P. Chankowski *et al.*, Nucl. Phys. **B417**, 101 (1994); D. Garcia and J. Sola, Mod. Phys. Lett. A **9**, 211 (1994); A. Dabelstein, W. Hollik, and W. Mosle, in *Perspectives for Electroweak Interactions in e^+e^- Collisions*, edited by B. A. Kniehl (World Scientific, Singapore, 1995), p. 345; D. Pierce *et al.*, Nucl. Phys. **B491**, 3 (1997).
- [22] H. T. Edwards, in *Annual Review of Nuclear and Particle Science*, edited by J. D. Jackson (Annual Reviews, Palo Alto, CA, 1985), Vol. 35, p. 605.
- [23] D0 Collaboration, S. Abachi *et al.*, Nucl. Instrum. Methods Phys. Res. A **338**, 185 (1994).
- [24] D0 Collaboration, M. Abolins *et al.*, Nucl. Instrum. Methods Phys. Res. A **280**, 36 (1989); D0 Collaboration, S. Abachi *et al.*, *ibid.* **324**, 53 (1993); D0 Collaboration, H. Aihara *et al.*, *ibid.* **325**, 393 (1993).
- [25] Q. Zhu, Ph.D. thesis, New York University, 1994, http://www-d0.fnal.gov/publications_talks/thesis/zhu/thesis_1side.ps.
- [26] T. C. Heuring, Ph.D. thesis, State University of New York at Stony Brook, 1993, http://www-d0.fnal.gov/publications_talks/thesis/heuring/thesis2s.ps.
- [27] J. W. T. McKinley, Ph.D. thesis, Michigan State University, 1996.
- [28] F. Carminati *et al.*, “GEANT Users Guide,” CERN Program Library W5013, 1991 (unpublished).
- [29] D0 Collaboration, S. Abachi *et al.*, Phys. Rev. D **52**, 4877 (1995).
- [30] G. A. Ladinsky and C. P. Yuan, Phys. Rev. D **50**, 4239 (1994).
- [31] P. B. Arnold and R. P. Kauffman, Nucl. Phys. **B349**, 381 (1991).
- [32] P. B. Arnold and M. H. Reno, Nucl. Phys. **B319**, 37 (1989); **B330**, 284E (1990).
- [33] J. Collins and D. Soper, Nucl. Phys. **B193**, 381 (1981); **B213**, 545E (1983); J. Collins, D. Soper, and G. Sterman, *ibid.* **B250**, 199 (1985).
- [34] G. Altarelli *et al.*, Nucl. Phys. **B246**, 12 (1984).
- [35] A. D. Martin, W. J. Stirling, and R. G. Roberts, Phys. Rev. D **50**, 6734 (1994); **51**, 4756 (1995).
- [36] D0 Collaboration, S. Abachi *et al.*, Phys. Rev. Lett. **75**, 1456 (1995).
- [37] G. Marchesini *et al.*, Comput. Phys. Commun. **67**, 465 (1992), release 5.7.
- [38] H. Plotow-Besch, Report No. CERN-PPE W5051, 1997, release 7.02.
- [39] E. Mirkes, Nucl. Phys. **B387**, 3 (1992).
- [40] J. Collins and D. Soper, Phys. Rev. D **16**, 2219 (1977).
- [41] F. A. Berends and R. Kleiss, Z. Phys. C **27**, 365 (1985); F. A. Berends *et al.*, *ibid.* **27**, 155 (1985).
- [42] H. L. Lai *et al.*, Phys. Rev. D **51**, 4763 (1995).
- [43] J. Botts *et al.*, Phys. Lett. B **304**, 159 (1993).
- [44] A. D. Martin, W. J. Stirling, and R. G. Roberts, Phys. Lett. B **306**, 145 (1993); **309**, 492E (1993).
- [45] T. Taylor Thomas, Ph.D. thesis, Northwestern University, 1997, http://www-d0.fnal.gov/publications_talks/thesis/thomas/thesis_2side.ps.
- [46] This test returns a confidence level κ that is uniformly distributed between 0 and 1 if the two spectra derive from the same parent distribution. See, e.g., A. G. Frodesen, O. Skjeggstad, and H. Tøfte, *Probability and Statistics in Particle Physics* (Columbia University Press, New York, 1979).

- [47] D. Casey, Ph.D. thesis, University of Rochester, 1997, http://www-d0.fnal.gov/publications_talks/thesis/casey/thesis.ps.
- [48] These unpublished fits were made available by the CTEQ Collaboration.
- [49] D0 Collaboration, S. Abachi *et al.*, Phys. Rev. Lett. **79**, 1203 (1997).
- [50] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **74**, 850 (1995).
- [51] U. Baur *et al.*, Phys. Rev. D **56**, 140 (1997); U. Baur *et al.*, in *Proceedings of the 1996 DPF/DPB Summer Study on High Energy Physics*, edited by D. G. Cassel, L. Trindle Gennari, and R. H. Siemann (Stanford Linear Accelerator Center, Stanford, 1997), Vol. 1, p. 514; U. Baur, S. Keller, and D. Wackerroth, *ibid.*, Vol. 1, p. 517; U. Baur, S. Keller, and W. K. Sakumoto, Phys. Rev. D **57**, 199 (1998).
- [52] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **65**, 2243 (1990); Phys. Rev. D **43**, 2070 (1991).
- [53] G. Degrassi *et al.*, Phys. Lett. B **418**, 209 (1998); G. Degrassi, P. Gambino, and A. Sirlin, Phys. Lett. B **394**, 188 (1997).